





# On the distribution and conservation status of some Mediterranean species new for the flora of Montenegro

Danijela Stešević<sup>1</sup> , Milica Stanišić-Vujačić<sup>1</sup> , Đorđije Milanović<sup>2</sup>  & Urban Šilc<sup>3</sup> 

**Key words:** alien species, conservation, IUCN assessment, new records.

**Ključne besede:** tujerodne vrste, ohranjanje, ocena IUCN, novi podatki.

Corresponding author:  
Danijela Stešević  
E-mail: danijela.stesevic@ucg.ac.me

Received: 13. 12. 2023  
Accepted: 30. 6. 2024



## Abstract

Investigations of dune habitats in Montenegro led to the discovery of 8 taxa new for the country: *Avellinia festucoides*, *Corynephorus divaricatus*, *Isoetes histrix*, *Juncus tenuis*, *Medicago doliata*, *Myosotis laxa* ssp. *cespitosa*, *Setaria parviflora*, and *Solenopsis laurentia* ssp. *gasparrinii*. Of particular importance are *Solenopsis laurentia* ssp. *gasparrinii* (not yet been recorded in the SE Adriatic) and *Isoetes histrix* (doubtfully reported for Albania). Since only one or small number of additional localities are expected in Montenegro, species are proposed as candidates for the national list of legally protected species: *Isoetes histrix*, *Solenopsis laurentia* ssp. *gasparrinii*, *Avellinia festucoides* and *Corynephorus divaricatus*. According to the IUCN criteria, the *Isoetes histrix*, *Solenopsis laurentia* ssp. *gasparrinii* and *Corynephorus divaricatus* are assessed as CR, while *Avellinia festucoides* as EN. Monitoring and eradication measures are proposed for alien species (*Setaria parviflora* and *Juncus tenuis*), especially for the second one, which is considered invasive.

## Izvleček

Med raziskavami peščenih sipin v Črni gori smo odkrili osem vrst in nižjih taksonov, ki so novi za Črno goro: *Avellinia festucoides*, *Corynephorus divaricatus*, *Isoetes histrix*, *Juncus tenuis*, *Medicago doliata*, *Myosotis laxa* ssp. *cespitosa*, *Setaria parviflora* in *Solenopsis laurentia* ssp. *gasparrinii*. Posebej pomembna sta *Solenopsis laurentia* ssp. *gasparrinii* (prvič zabeležena na JV delu Jadrana) in *Isoetes histrix* (le dvomljive navedbe za Albanijo). Sledeče taksone smo predlagali za uvrstitve na nacionalni seznam pravno zavarovanih vrst, saj lahko v Črni gori pričakujemo le majhno število dodatnih nahajališč: *Isoetes histrix*, *Solenopsis laurentia* ssp. *gasparrinii*, *Avellinia festucoides* in *Corynephorus divaricatus*. V skladu z merili IUCN smo taksone *Isoetes histrix*, *Solenopsis laurentia* ssp. *gasparrinii* in *Corynephorus divaricatus* uvrstili v kategorijo CR, vrsto *Avellinia festucoides* pa v EN. Za tujerodne vrste (*Setaria parviflora* in *Juncus tenuis*) predlagamo redni monitoring in ukrepe za odstranjevanje, še posebej za drugo vrsto, ki jo v nekaterih državah obravnavajo kot invazivno.

1 University of Montenegro, Faculty of Natural Sciences and Mathematics, Podgorica, Montenegro.

2 University in Banja Luka, Faculty of Forestry, Banja Luka, Bosnia and Herzegovina.

3 Research Centre of the Slovenian Academy of Sciences and Arts (ZRC SAZU), Jovan Hadži Institute of Biology, Ljubljana, Slovenia.

## Introduction

The Velika plaža area in Ulcinj and its hinterland is considered one of the most botanically researched sites on the southeastern Adriatic, with its flora, vegetation and habitats documented by numerous publications (Rohlena, 1942; Bubanja et al., 2019; Stešević et al., 2020; Caković et al., 2021) or herbarium specimens in e.g. TGU, NHMM, BEO, GZU, PRC (Thiers, 2024). In the dune systems of Velika plaža and its hinterland, ca. 970 taxa of vascular plants have been recorded so far (Bubanja et al., 2019; Stešević et al., 2020; Stešević et al., 2021; Caković et al., 2021), and ca. 30 plant communities have been listed (Blečić & Lakušić, 1976; Mijović, 1994; Stešević et al., 2020; Milanović et al., 2021). This area is characterised by a mosaic of different coastal, dune and riparian habitats that have high conservation value, but are also subject to strong and constant anthropogenic pressures that led to degradation or complete disappearance of the most sensitive habitats (Šilc et al., 2020; Milanović et al., 2021). This unique area in Montenegro, still does not enjoy an adequate protection status in the country. So far, only a part of the dunes on Velika plaža and the Ulcinj Salina have the status of a protected area, more precisely, the first one is a natural monument, category III and IUCN category of protected area III, and the second one is a nature park, category II and IUCN category of protected area IV (CGIS, 2017). However, for the former

area, the boundary has not yet been officially established, so a revision is needed.

With the primary goal of further vulnerability monitoring of NATURA 2000 habitat types to invasion by alien plant species (Stešević et al., 2017; Šilc et al., 2019), a systematic survey of the Velika plaža hinterland was conducted in 2021. Two new species for the adventitious flora of Montenegro were found, but also some peculiarities for the native flora were recorded, such as *Isoetes hirtix* and *Solenopsis laurentia* ssp. *gasparrinii*. Other new records for the flora of this place are an incentive for further research (synecological, vegetation, conservation, etc.), but also for monitoring of priority species, either from the point of view of protection or control of dispersal. By highlighting the biological value of the Velika plaža and its hinterland and also drawing attention to the threats that lead to its further devastation and the disappearance of important habitats and then stenobiontic species, we contribute to the protection of this area.

## Material and methods

The field survey was conducted in 2021 in the hinterland of Velika plaža near Ulcinj (Figure 1). With the length of ca. 12 km, Velika plaža is considered to be the longest beach on the eastern Adriatic coast, which despite the anthropogenic pressure still has well-developed vegetation (Šilc et al., 2016). Due to the fact that 11 NATURA 2000



**Figure 1:** Geographic position of Velika plaža near Ulcinj (Montenegro).  
**Slika 1:** Geografska lokacija Velike plaže pri Ulcinju (Črna gora).

habitats are reported for the beach and its hinterland (Stešević et al., 2020; Milanović et al., 2021), an initial goal of the survey was to monitor plant invasions in following non-forest NATURA 2000 habitats: 2240 *Brachypodietalia* dune grasslands with annuals, 3170 \*Mediterranean temporary ponds, 6220\* Pseudo-steppe with grasses and annuals of the *Thero-Brachypodietea*, 6420 Mediterranean tall humid herb grasslands of the *Molinio-Holoschoenion*, and 6540 Sub-Mediterranean grasslands of the *Molinio-Hordeion secalini*.

The collected plant material has been deposited in the Herbarium Collection of the University of Montenegro (TGU) and in the Herbarium of the Faculty of Forestry of the University of Banja Luka (BALU).

Specimens of *Solenopsis* were identified according to Brullo et al. (2023), of the *Isoetes* according to Bagella et al. (2015) and Troia & Greuter (2014, 2015) and *Juncus tenuis* according to Verloove (2010), while the specimens of the other species were identified according to Tutin et al. (1972, 1976, 1980), Pignatti (1982), and Nikolić (2020). The nomenclature of the species follows POWO (2023), except for *Solenopsis* (Brullo et al., 2023).

The assessment of threats to native species on national/regional level was carried out according to the IUCN method (IUCN Category Standards and Petitions Committee, 2022). The spatial analysis sensu calculation of the geographic range size of a species (EOO – the extent of occurrence) and the area in which a species occurs (AOO – the area of occupancy) for the Red List assessment was carried out using GeoCAT – Geospatial Conservation Assessment Tool algorithms (Bachman et al., 2011). For the final regional assessment (Step 3) the potential rescue effects from populations occurring outside of the assessment region were evaluated and corresponding up- or down-listing of the species' extinction risk status at the regional level was considered.

## Results and discussion

*Isoetes histrix* Bory, Montenegro, Ulcinj – Velika plaža, hinterland, temporary Mediterranean ponds, 07.05.2021, 19.05.2021, 02.6.2021, Det. D. Stešević, M. Stanišić-Vujačić, Đ. Milanović (Herbarium TGU, BALU).

In the ecological environment of the Velika plaža hinterland, this summer-deciduous perennial species spends part of its annual cycle under water and begins the emergent life phase from mid or late March. The above-ground parts of the plant (Figure 6a) remain until the beginning of the summer drought, which starts as early as mid-June. Compared to other taxa from the complex *Isoetes histrix* s.l., the species in the narrower sense (*Isoetes histrix* s.str.) is characterized by persistent black phyllopodia on

top of the corm and around the leaf bases (Figure 6B), longer than 3 mm, while the megaspores are tuberculate and measure up to 500 µm. In addition to individuals with long phyllopodium prickles, those with shorter ones were also collected, but due to the impossibility of a more precise comparison of spores, observation of microspore and megaspore ornamentation using SEM, the question of whether this is only an infraspecies variation or the co-existence of different species remains open. Compared to other species of this genus, the species *I. histrix* is characterized by exceptional morphological variability in terms of leaf length, phyllopodia, and spore ornamentation (Bagella et al., 2011). According to Troia & Greuter (2014), the species complex *I. histrix* s.l. consists of two taxa, and according to Bagella et al. (2015) of three. Namely, the second group of authors separates *I. gymnocarpa* from *I. sicula*, while the first group of authors synonymizes them. Recently, Romanov et al. (2024) recorded the taxon *I. gymnocarpa* (= *I. sicula*) near the Ulcinj Salina, which is an additional reason for further research not only on taxon diversity but also on its synecology.

Specimens of *Isoetes histrix* s.str. were collected on 3170\* – Mediterranean temporary ponds (Figure 3), where it grows together with *Polypogon maritimus*, *Trifolium lappaceum*, *Isolepis cernua*, *Juncus bufonius*, *Juncus capitatus*, *Leontodon saxatilis*, *Trifolium patens*, *Lotus angustissimus*, *Linum bienne*, *Poa jubata*, *Plantago lanceolata*, *Avellinia festucoides*, *Radiola linoides*.

Seasonally flooded habitats, including Mediterranean temporary ponds, are common habitats for species included into *Isoetes histrix* s.l. (Biondi et al., 2012; Bagella & Podani, 2017; Tomaselli et al., 2022). It is not uncommon that two taxa from the complex grow together, for example, *I. histrix* and *I. sicula* in plant communities *Solenopsidetum caespitosae* and *Poa jubatae-Isoetetum histricis* (Tomaselli et al., 2022). Therefore, it is important to resolve the identity of the species of the genus *Isoetes* inhabiting Mediterranean temporary ponds in Montenegro, and then to determine the plant communities in which they thrive.

Data on the distribution of *Isoetes histrix* in the eastern part of the Adriatic Sea are rather scarce. So far, the species has not been recorded in neighbouring Croatia (Nikolić, 2023), while in Albania it is considered “occurrence not proven” - i.e. it is included in the checklist because the species is mentioned in some synthetic works, but without specifying the exact locality or supporting it with a herbarium voucher (Barina et al., 2018).

In the territory of Montenegro, species of the genus *Isoetes* have been intensively searched for in the coastal area of Lake Skadar in the last decade, as this area is considered a potential habitat for its amphibious and terres-

trial representatives (Hadžiablahović, pers. comm.). Recent findings confirmed two new taxa (both from *Isoetes histrix* complex) for the lycopod flora in Montenegro.

In terms of conservation importance, *I. histrix* is listed as Least Concern in the European Red List of Species (European Environmental Agency, 2023), as Least Concern in the 2016 IUCN Red List of Threatened Species (Christenhusz et al., 2017), in England it has the status VU (Stroh et al., 2014), in France LC, and in certain regions of France: EN (Provence-Alpes-Côte d’Azur, Poitou-Charentes), CR (la région Centre), VU (d’Aquitaine, Pays de la Loire), NT (Bretagne) – Inventaire national du patrimoine naturel (2023).

Despite the current uncertainty regarding the intra-population variability of *Isoetes histrix* in the Velika plaža hinterland or coexistence with another taxon from the species

complex of the same name, we consider *I. histrix* a good candidate for the national list of legally protected plants. A final assessment of the IUCN category still requires further research. Based on available data (Table 1, Figure 2), the species is preliminary assessed as CR, according to criteria B1ab(ii,iii,v). When assessing the regional population and status in neighbouring countries, we conclude that no changes are needed compared to step 2, so the preliminary category remains unchanged. The main threat is the change in the ecosystem due to urban sprawl, shrub encroachment, but also due to changes in the climate and water regime.

*Solenopsis laurentia* ssp. *gasparrinii* (Tineo) Brullo, Montenegro, Ulcinj – Velika plaža, hinterland Mediterranean temporary ponds, 21. 05. 2021, 03. 6. 2021, Det. D. Stešević, M. Stanišić-Vujačić, Đ. Milanović (Herbarium TGU, BALU).

**Table 1:** Assessment of *Isoetes histrix*, *Solenopsis laurentia* ssp. *gasparrinii*, *Avellinia festuroides* and *Corynephorus divaricatus* to IUCN criteria.

**Tabela 1:** Ocena ogroženosti po merilih IUCN za taksone *Isoetes histrix*, *Solenopsis laurentia* ssp. *gasparrinii*, *Avellinia festuroides* in *Corynephorus divaricatus*.

Species	EOO (km <sup>2</sup> ) /assessment	AOO (km <sup>2</sup> ) /assessment	Assessment category
<i>Isoetes histrix</i>	3.733 (CR)	24 (EN)	CR B1ab(ii,iii,v)
<i>Solenopsis laurentia</i> ssp. <i>gasparrinii</i>	0.003 (CR)	8 (CR)	CR B1+2ab(ii,iii,v)
<i>Avellinia festuroides</i>	5.026 (CR)	24 (EN)	EN B1ab(ii,iii,v)
<i>Corynephorus divaricatus</i>	2.336 (CR)	24 (EN)	CR B1ab(ii,iii,v)



**Figure 2:** Locations of *Isoetes histrix* s.l. localities in the Velika plaža hinterland.

**Slika 2:** Nahajališča vrste *Isoetes histrix* s.l. v zaledju Velike plaže.


Temporarily flooded and grassy habitats in the Velika plaža hinterland are inhabited by *Solenopsis laurentia* ssp. *gasparrinii* (Tineo) Brullo, *Myosotis laxa* ssp. *cespitosa* (Schultz) Nordh. and *Avellinia festucoides* (Link) Valdés & H. Scholz (Figure 6d, e, g Figure 3, Figure 4). The finding of the former taxon is particularly important, because it has not been recorded before in the Balkan Peninsula (Brullo et al., 2023). The genus *Solenopsis* is also new for the territory of Montenegro (Rohlena, 1942; Pulević, 2005).

It is a species complex with several taxa at specific and infraspecific ranks. The genus *Solenopsis* is considered endemic to the Mediterranean region (Crespi et al., 1998). Of the *S. laurentia* subordinate taxa, the *S. laurentia* ssp. *laurentia* has the widest distribution – throughout the Mediterranean region. The subspecies *gasparrinii* occurs in Sicily, Apulia, Sardinia, Corsica, and northern Morocco (Brullo et al., 2023). However, Montenegro has now been included in its distribution range. So far, representatives of the genus *Solenopsis* have not been recorded in neighbouring Croatia (Nikolić, 2023), while the review list of the Flora of Albania mentions *Solenopsis laurentia*, but as “occurrence not proven” (Barina et al., 2018).

In the hinterland of Velika plaža in Ulcinj, *Solenopsis laurentia* ssp. *gasparrinii* was detected in Mediterranean

temporary ponds together with *Polygonum maritimum*, *Carex flacca* ssp. *erythrostachys*, *Cyperus flavescens*, *Isolepis cernua*, *Lotus angustissimus*, *Trifolium lappaceum*, *Juncus bufonius*, *Myosotis laxa* ssp. *cespitosa*. The life cycle begins in April, when the water has already receded but the soil is still quite marshy. The cycle ends in late May or early June, when the summer drought begins. In the whole distribution range this taxon is closely associated with the Mediterranean temporary ponds (Brullo et al., 2023).

In Italy, the conservation status of the taxon is Endangered [EN, B2ab(ii,iii)] (Brullo et al., 2023). Taking into account the ecological connection of the species to Mediterranean temporary ponds, which so far have been recorded only in the hinterland of Velika plaža in Ulcinj, it is highly unlikely to expect the species elsewhere in Montenegro. According to the estimated number of locations, EOO and AOO (Table 1), the species is provisionally assessed as CR according to IUCN criteria B1+2ab(ii,iii,v). When assessing the regional population and status in neighbouring countries, we conclude that no changes are necessary compared to step 2, so the provisional category remains unchanged. Due to the relatively high anthropogenic pressure in the area, especially urban sprawl, the range (ii), habitat quality (iii), and number of mature individuals (v) are predicted to steadily decline.



**Figure 3:** Distribution of *Juncus tenuis*, *Medicago doliata*, *Myosotis laxa* ssp. *cespitosa*, *Setaria parviflora*, *Solenopsis laurentia* ssp. *gasparrinii* in the hinterland of Velika plaža. Numbers in the bracket represent the number of sites on which species were found.

**Slika 3:** Razširjenost taksonov *Juncus tenuis*, *Medicago doliata*, *Myosotis laxa* ssp. *cespitosa*, *Setaria parviflora*, *Solenopsis laurentia* ssp. *gasparrinii* v zaledju Velike plaže. Številke v oklepajih predstavljajo število nahajališč, kjer smo našli posamezen takson.

*Medicago doliata* Carmign. (syn. *Medicago aculeata* Willd.), Montenegro, Ulcinj – Velika plaža, hinterland, dry grassland, 21. 05. 2021, 21. 06. 2021, Det. D. Stešević (Herbarium TGU).

*Medicago doliata* was found on the dry grassland (Figure 3) dominated by the species *Aegilops neglecta* with the following species also present: *Crepis neglecta*, *Lagurus ovatus*, *Poa bulbosa*, *Vulpia ciliata*, *Onobrychis caput-galli*, *Plantago bellardi*, *Trifolium cherleri*, *Trifolium scabrum*, *Sherardia arvensis*, and *Catapodium rigidum*. It differs from other species of this genus recorded in the hinterland of Velika plaža (Bubanja et al., 2019) by its globose to ellipsoid legumes with dense, short, sometimes glandular hairs and mostly conical and straight spines (Figure 6c). The natural range includes dry grassland habitats in the Mediterranean region and Iran (Pignatti, 1982; POWO, 2023).

It has the status LC at the global level and in Europe (European Environmental Agency, 2023). It is expected that the species will be recorded in more localities during further surveys of the coastal area of Montenegro. According to current information it is not possible to adequately assess the threat status, thus we propose DD (Data Deficient).

*Myosotis laxa* ssp. *cespitosa* (syn. *Myosotis cespitosa* Schultz, *Myosotis laxa* var. *cespitosa* (Schultz) Apelgren, *Myosotis scorpiooides* ssp. *cespitosa* (Schultz) F. Hermann), Montenegro, Ulcinj – Velika plaža, hinterland, wet meadows, 15. 05. 2021, 08. 06. 2021; Mediterranean temporary ponds, 21. 05. 2021, Det. D. Stešević (Herbarium TGU). In addition to Mediterranean temporary ponds, it was also detected in wet meadows (Habitat 6540), where it grows together with *Alopecurus rendlei*, *Ranunculus sardous*, *Trifolium resupinatum*, *Anacamptys laxiflora*, *Silene flos-cuculi*, *Lotus angustissimus*, *Poa trivialis* (Figure 1). From the other species of the genus *Myosotis*, which have been recorded so far in the hinterland of Velika plaža: *Myosotis arvensis*, *M. discolor* (Bubanja et al., 2019) and *M. ramosissima* (TGU 2155908), it differs by the straight hairiness of the calyx (Figure 6g), while from the *M. scorpiooides* that is common species in wet habitats in Montenegro (Rohlena, 1942), by its annual form. Due to the characteristic that the calyx remains open on the fruit, it is possible that it is confused with *M. ramosissima* or *M. arvensis* if identified on pressed herbarium material.

Considering the size of the range – Temperate Eurasia and NW Africa (POWO, 2023), the ecology of the species and its attachment to a wide number of wet habitats, banks of standing and flowing waters, wet meadow ditches, edges of floodplain forests, sedge stands, Mediterranean temporary wetland systems (Grau & Merxmüller,

1972; Pignatti, 2005; Gigante et al., 2013), it is realistic to expect that this taxon is present in a larger number of localities in Montenegro.

At the European level, the species has no conservation status and has not yet been assessed for the IUCN Red List (European Environmental Agency, 2023). However, it is Red Listed in some countries (e.g. Czech Red List, Grulich, 2017). As in the case of the previous taxon, the proposed category for threat status in Montenegro is DD (Data Deficient).

*Avellinia festucooides* (Link) Valdés & H. Scholz (syn. *Avellinia michelii* (Savi) Parl.), Montenegro, Ulcinj – Velika plaža, hinterland, Mediterranean temporary pond 03. 06. 2021, dry grasslands 17. 06. 2021, Det. D. Stešević, Đ. Milanović (Herbarium TGU, BALU).

In the Velika plaža hinterland (Figure 4), this grass was recorded in rather different habitats, such as Mediterranean temporary ponds (Habitat 3170), Humid dune slack (Habitat 2190), in fringe vegetation of *Scirpoides holoschoenus* community, in Wooded dunes with *Pinus pinea* and/or *Pinus pinaster* (Habitat \*2270), *Brachypodietalia* dune grasslands with annuals (Habitat 2240) and *Vulpio-Lotion* annual grasslands (Habitat \*6220). It belongs to a genus that has not yet been recorded in the flora of Montenegro, while in the past it has been recorded in Croatia on the Dalmatian islands of Biševo and Vis in numerous populations on dry, sunny and grassy sandy habitats, as well as sporadically in garrigue and along roads (Pavletić, 1972). This Mediterranean species (POWO, 2023) prefers dry, open, sandy habitats, usually near the sea (Tutin, 1976; Pignatti, 1982). It is characterized by a delicate annual herbaceous form, a compact panicle with multi-flowered spikelets (Figure 6d), glumes very unequal, the upper of which is almost as long as the floret, the lemma 2-pinnate at the apex, with a short awn from the sinus, the palea shorter than the lemma, deeply 2-feed. We assume that the discovery of the species was postponed for a long time due to a possible confusion with similar species of the genera *Trisetaria* and *Koeleria*.

At the European level, the species has no protection status and it has not yet been assessed for the IUCN Red List (European Environmental Agency, 2023). In Croatia, this rare grass species has been assigned the status NT (Nikolić, 2023).

Due to restricted distribution, the species presents a good candidate for the national list of legally protected species. The sandy and maritime habitats, which are considered as ecological preference of the species (Tutin, 1976; Pignatti, 1982), are found on Velika plaža, its hinterland and a few small patches further along the coast. In a recently conducted survey of the patches, the species



Figure 4: Distribution on *Avellinia festucoides* in Montenegro, in the hinterland of Velika plaža.

Slika 4: Razširjenost vrste *Avellinia festucoides* v Črni gori v zaledju Velike plaže.

was not recorded. According to the estimated number of locations, EOO and AOO (Table 1), the species is provisionally classified as CR according to IUCN criteria B1ab(ii,iii,v). The assessment of the regional population and status in neighbouring countries in the final assessment category is a downgrade from step 2 in EN B1ab(ii,iii,v). Taking into account the relatively high anthropogenic pressure on the area, especially urban sprawl, a steady decline in range (ii), habitat quality (iii) and number of mature individuals (v) is predicted.

*Corynephorus divaricatus* (Pourr.) Breistr., Montenegro, Ulcinj – Velika plaža hinterland, dune grassland 15. 05. 2021, 02. 06. 2021, 08. 06. 2021, Det. D. Stešević (Herbarium TGU, BALU).

It is common on dry, sandy and decalcified grasslands, both on *Brachypodietalia* dune grasslands with annuals (Habitat 2240) and *Vulpio-Lotion* annual grasslands (Habitat \*6220), where it grows with relatively low cover of up to 5%. It is characterized by an annual, erect, and rather graceful form, a loose panicle whose two-flowered panicles contain a short, clavate awn strongly enlarge towards apex (Figure 6f), articulated near the middle, and with a ring of short hairs at the joint. Although it is not explicitly mentioned as part of the flora of Montenegro, in the overview of plant communities of Montenegro (Blečić & Lakušić, 1976) the association *Laguro-Corynephorum divaricati* H-ić 1964 is mentioned for the coastal part of

the country. So far, this community has not been documented, either in the published references or in the manuscripts. According to our observations, *Corynephorus divaricatus* is common in the hinterland of Velika plaža (Figure 5), but never with high abundance and forming the communities that could be classified into mentioned association. This grass is a species of Mediterranean range (POWO, 2023), characteristic of pastures and grasslands (Pignatti, 1982), so the habitat in the hinterland of Velika plaža can be considered as a typical one. As for the conservation status, it has no protection status at European level and has not yet been assessed for the IUCN Red List (European Environmental Agency, 2023). In Italy it has the status LC (Orsenigo et al., 2020), and in Croatia CR (Nikolić, 2023). The species can be considered a good candidate for the national list of legally protected species.

As in the case of the *Avellinia festucoides*, the *C. divaricatus* was not recorded elsewhere along the coast, thus Velika plaža with the hinterland is considered the only site in Montenegro. According to the estimated number of location, EOO and AOO (Table 1), the species is preliminary assessed as CR according to IUCN criteria B1ab(ii,iii,v). In the assessment of the regional population and status in neighbouring countries, the preliminary category is left unchanged. Taking into account a relatively high anthropogenic pressure on the area, especially urban sprawl, a steady decline in range (ii), habitat quality (iii) and number of mature individuals (v) is predicted.



Figure 5: Distribution of *Corynephorus divaricatus* in the hinterland of Velika plaža.

Slika 5: Razširjenost vrste *Corynephorus divaricatus* v zaledju Velike plaže.

*Juncus tenuis* Willd., Montenegro, Ulcinj – Velika plaža, hinterland, wet sites, 17. 06. 2021, Det. D. Stešević (Herbarium TGU), Figure 6h.

Sand embankments between artificial water bodies in the hinterland of Velika plaža were colonized by *Juncus tenuis*, *Fimbristylis dichotoma*, *Cyperus fuscus* and *Eleocharis acicularis* (Figure 3). *Juncus tenuis* s.str. is distinguished from other species in the *Juncus tenuis* complex by its long leaf sheath auricles (Verloove, 2010). This North American species has a secondary distribution across most of Europe (POWO, 2023), where it inhabits a wide range of habitats: dry, sandy, and more acidic soils, tracks or clearings in forests, disturbed heaths, along rivers, and other wet or urban habitats (Verloove, 2010). Seeds are well adapted to human or animal dispersal and mud colonization, so it is common along trails and roads (Salisbury, 1974).


It is on the global list of invasive species (Global Invasive Species Database, 2023), but not on the EU list. This North American species was in Europe first recorded in the UK in 1790 (Arianoutsou et al., 2021). Nevertheless, it has the status of invasive species in neighbouring countries (Croatia – Nikolić, 2023; Serbia – Lazarević et al., 2012; Bosnia and Herzegovina – Maslo, 2016), while it is extinct casual in Albania (Barina et al., 2014). Further monitoring of this species is needed to determine its status. In future research, attention should also be paid

to other species from the complex, such as *J. dudleyi*, which was recently recorded in Bosnia and Herzegovina (Milanović et al., 2018).

*Setaria parviflora* (Poir.) Kerguelen, Montenegro, Ulcinj – Velika plaža hinterland, ruderalized lawn 15. 07. 2021, Det. D. Stešević, Đ. Milanović (Herbarium TGU, BALU).

On the ruderalized lawn dominated by annual species (e.g. *Avena barbata*, *Vulpia ligustica*, *Haynaldia villosa*, *Lagurus ovatus*, *Vicia grandiflora*), a perennial plant from the genus *Setaria* – *S. parviflora* – was recorded (Figure 3). It differs from the species of this genus previously recorded in the Velika plaža hinterland (*S. viridis*, Bubanja et al., 2019) by its perennial form, spike-like panicles up to 10 cm long and numerous yellow to purple bristles up to 6 times longer than the spikelets (Hitchcock, 1971). This American species has a secondary distribution in the Mediterranean region, southern Africa and Madagascar, Southeast Asia, and Polynesia (POWO, 2023). Although it was first recorded in Europe more than two centuries ago (1800 in Austria, Arianoutsou et al., 2021), it has not spread widely. In Italy and Croatia, it has the status of a naturalized neophyte (Galasso et al., 2018; Nikolić, 2023). Further monitoring of this species is needed to determine its status, either as a casual alien, naturalized neophyte or invasive species.





**Figure 6 (Slika 6):** A, B – *Isoetes histrix*, C – *Medicago doliata*, D – *Avellinia festucoides*, E – *Solenopsis laurentia* ssp. *gasparrinii*, F – *Corynephorus divaricatus*, G – *Myosotis laxa* ssp. *cespitosa*, H – *Juncus tenuis*.

## Conclusions


Our study and results support that it would be justified to propose the extension of the protected area of sand dunes to the hinterland itself, where very rare plant species and also NATURA habitats are found, such as: 1150 Coastal lagoons, 1410 Mediterranean salt meadows (*Juncetalia maritimi*), 2190 Humid dune slacks, 3170 \*Mediterranean temporary ponds and 6420 Mediterranean tall humid herb grasslands of the *Molinio-Holoschoenion* (Milanović et al., 2021). It would be worthwhile to combine sand dunes and hinterland into one protected area with a buffer zone due to exceptional species and vegetation types. Nevertheless, the human impact has been very destructive in the last decades by urbanisation and tourism, therefore the first step would be to at least protect the most valuable and preserved parts of Velika plaža hinterland.


## Acknowledgements

The authors would like to thank Angelo Troia and Simonetta Bagella for valuable discussions considering *Isoetes histrix* s.l. and confirmation of identification of *Isoetes histrix* s.str.; Gianmarco Tavilla and Salvatore Brullo for confirmation of identification of *Solenopsis laurentia* ssp. *gasparrinii*; Peter Glasnović for valuable advice about IUCN criteria and categories; Iztok Sajko for preparing the maps; the anonymous reviewers for kind suggestions for the improvement of the manuscript. Urban Šilc was supported by ARIS grant P1-0236.

### ORCID iDs

Danijela Stešević  <https://orcid.org/0000-0003-0015-7141>

Milica Stanišić-Vujačić  <https://orcid.org/0000-0002-5711-8275>

Đorđije Milanović  <https://orcid.org/0000-0002-9130-7600>

Urban Šilc  <https://orcid.org/0000-0002-3052-699X>

## References

Arianoutsou, M., Bazos, I., Christopoulou, A., Kokkoris, Y., Zikos, A., Zervou, S., Delipetrou, P., Cardoso, A. C., Deriu, I., Gervasini, E., & Tsiamis, K. (2021). Alien plants of Europe: introduction pathways, gateways and time trends. *PeerJ*, 9, e11270. <https://doi.org/10.7717/peerj.11270>

Bachman, S., Moat, J., Hill, A. W., de la Torre, J., & Scott, B. (2011). Supporting Red List threat assessments with GeoCAT: geospatial conservation assessment tool. *ZooKeys*, 150, 117–126. <https://doi.org/10.3897/zookeys.150.2109>

Bagella, S., Peruzzi, L., Caria, M. C., & Filigheddu, R. (2015). Unravelling the taxonomy and nomenclature of the *Isoetes histrix* Bory species complex (Isoetaceae, Lycopodiidae). *Turkish Journal of Botany*, 39(2), 383–387. <https://doi.org/10.3906/bot-1404-121>

Bagella, S., Podani, J. (2017). A large-scale assessment of *Isoetes histrix* s.l. swards in the Mediterranean basin. *Plant Sociology*, 54(1), 129–136. <https://doi.org/10.7338/pls2017541/06>

Barina, Z., Rakaj, M., Somogyi, G., Erős-Honti, Z., & Pifkó, D. (2014). The alien flora of Albania: history, current status and future trends. *Weed Research*, 54(2), 196–215. doi: 10.1111/wre.12061

Barina, Z., Somogyi, G., Pifkó, D., Rakaj, M. (2018). Checklist of vascular plants of Albania. *Phytotaxa* 378, 1–339. <https://doi.org/10.11646/phytotaxa.378.1.1>

Biondi, E., Burrascano, S., Casavecchia, S., Copiz, R., Del Vico, E., Galdenzi, D., Gigante, D., Lasen, C., Spampinato, G., Venanzoni, R., Zivkovic, L., & Blasi, C. (2012). Diagnosis and syntaxonomic interpretation of Annex I Habitats (Dir. 92/43/EEC) in Italy at the alliance level. *Plant Sociology*, 49(1), 5–37. <https://doi.org/10.7338/pls2012491/01>

Blečić, V., & Lakušić, R. (1976). List of plant communities of Montenegro. *Glasnik Republičkog Zavoda za Zaštitu Prirode i Privrednjackog Muzeja u Titogradu*, 9, 57–98 (in Montenegrin).

Brullo, S., Brullo, C., Cambria, S., Minissale, P., Sciandrello, S., Tavilla, G., Siracusa, G., Tomaselli, V., & Giusso del Galdo, G. (2023). Taxonomical remarks on *Solenopsis laurentia* (Campanulaceae) in Italy. *Phytotaxa*, 584(2), 59–88. <https://doi.org/10.11646/phytotaxa.584.2.1>

Bubanja, N., Šinžar-Sekulić, J., & Stevanović, V. (2019). Contribution to the flora of Velika plaža and its vicinity in Ulcinj (Montenegro). *Biologia Serbica*, 41(1), 13–44. <https://doi.org/10.5281/zenodo.3263954>

Caković, D., Stešević, D., & Vuksanović, S. (2021). *Pulicaria sicula* (L.) Moris, a new species in the flora of Montenegro and the westernmost record in the Balkan Peninsula. *Ecologica Montenegrina*. 43, 51–55. <https://doi.org/10.37828/em.2021.43.7>

CGIS (2017). Web portal za nacionalno zaštićena područja, <http://prirodainfo.me/>, accessed July 2023.

Christenhusz, M., Bento Elias, R., Dyer, R., Ivanenko, Y., Rouhan, G., Rumsey, F., & Väre, H. (2017). *Isoetes histrix* (Europe assessment). *The IUCN Red List of Threatened Species 2017*, e.T164366A85431088. Accessed on 18 August 2023.

Crespo, MB, Serra, L. & Juan, A. (1998). *Solenopsis* (Lobeliaceae): a genus endemic in the Mediterranean Region. *Plant Systematic and Evolution*, 210, 211–229. <https://doi.org/10.1007/BF00985669>

European Environmental Agency (2023, August 16). European Red list of species. <https://www.eea.europa.eu/en/datahub/datahubitem-view/202f3c2e-54a9-4ff4-a1da-ed/ca524f634>

Galasso, G., Conti, F., Peruzzi, L., Ardenghi, NMG, Banfi, E., Celesti-Grappow, L., Albano, A., Alessandrini, A., Bacchetta, G., Ballelli, S., Bandini, Mazzanti, M., Barberis, G., Bernardo, L., Blasi, C., Bouvet, D., Bovio, M., Cecchi, L., Del Guacchio, E., Domina, G., Fascetti, S., Gallo, L., Gubellini, L., Guiggi, A., Iamonic, D., Iberite, M., Jiménez-Mejías, R., Lattanzi, E., Marchetti, D., Martinetto, E., Masin, RR, Medagli, P., Pas-salacqua, NG, Peccenini, S., Pennesi, R., Pierini, B., Podda, L., Poldini, L., Prosser, F., Raimondo, FM, Roma-Marzio, F., Rosati, L., Santangelo, A., Scoppola, A., Scortegagna, S., Selvaggi, A., Selvi, F., Soldano, A., Stinca, A., Wagensommer, RP, Wilhalm, T, Bartolucci, F. (2018). An updated checklist of the vascular flora alien to Italy. *Plant Biosystems*, 152(3), 556–592. <https://doi.org/10.1080/11263504.2018.1441197>

Gigante, D., Maneli, F., & Venanzoni R. (2013). Mediterranean temporary wet systems in inland Central Italy: ecological and phytosociological features. *Plant Sociology*, 50(2), 93–112. <https://doi.org/10.7338/pls2013502/06>

- Global Invasive Species Database (2023). Species profile: *Juncus tenuis*. <http://www.iucngisd.org/gisd/speciesname/Juncus+tenuis> on 08-09-2023.
- Grau, J. & Merxmüller, H. (1972). *Myosotis*. In T.G. Tutin, et al. (Eds.), *Flora Europaea* (3, pp. 111–117), Cambridge University Press.
- Grulich V. (2017). Červený seznam cévnatých rostlin ČR [The Red List of vascular plants of the Czech Republic]. *Příroda*, 35, 75–132.
- Hitchcock, A. S. (1971). *Manual of Grasses of the United States*. Courier Corporation.
- Inventaire national du patrimoine naturel – INPN (2023, October 13). *Isoetes histrix*. [https://inpn.mnhn.fr/espece/cd\\_nom/103842/tab/statut](https://inpn.mnhn.fr/espece/cd_nom/103842/tab/statut)
- IUCN Standards and Petitions Committee (2022). Guidelines for Using the IUCN Red List Categories and Criteria. Version 15.1. <https://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Lazarević, P, Stojanović, V, Jelić, I, Perić, R, Krsteski, B, Ajtić, R, Sekulić, N, Branković, S, Sekulić, G, Bjedov, V (2012). A preliminary list of invasive species in Serbia, with general measures of control and reduction as a basis of future legal acts. *Zaštita prirode*, 62(1), 5–31 (in Serbian).
- Maslo, S. (2016). Preliminary list of invasive alien plant species (IAS) in Bosnia and Herzegovina. *Herbologia*, 16(1), 1–14.
- Mijović, A. (1994). The xerohalophytic sand communities of the Long beach. *Glasnik Instituta za Botaniku i Botaničke Bašte Univerziteta u Beogradu*, 28, 147–157.
- Milanović, Đ, Maslo, S., & Šarić, Š. (2018). Four neophytes new for the flora of Bosnia and Herzegovina. *Botanica Serbica*, 42(1), 139–146. <https://doi.org/10.5281/zenodo.1173570>
- Milanović, Đ, Caković, D, Hadžiablahović, S, Vuksanović, S, Mačić, V, Stešević, D, Stanišić-Vujačić, M, Biberdžić, V, Lakušić, D. (2021). Manual for identification of habitat types of Montenegro of importance for the European Union with processed main indicator species. Agencija za zaštitu životne sredine Crne Gore, Univerzitet u Banjoj Luci – Šumarski fakultet (in Montenegrin).
- Nikolić, T. (2020). *Flora Croatica*. Alfa.
- Nikolić T. (Ed.) (2023). Flora Croatica Database. Faculty of Science, University of Zagreb. <http://hirc.botanic.hr/fcd>
- Orsenigo, S, Fenu, G, Gargano, D, Montagnani, C, Abeli, T, Alessandrini, A, Bacchetta, G, Bartolucci, F, Carta, A, Castello, M, Cogoni, D, Conti, F, Domina, G, Foggi, B, Gennai, M, Gigante, D, Iberite, M, Peruzzi, L, Pinna, MS, Prosser, F, Santangelo, A, Selvaggi, A, Stinca, A, Villani, M, Wagensommer, RP, Tartaglioni, N, Duprè, E, Blasi, C, & Rossi, G. (2020). Red List of threatened vascular plants in Italy. *Plant Biosystems*, 155(2), 310–335. <https://doi.org/10.1080/11263504.2020.1739165>
- Pavletić, Z. (1972). *Avellinia michelii* (Savi) Pari. — nova biljka u flori Jugoslavije. *Acta Botanica Croatica*, 31(1), 211–213. <https://hrack.srce.hr/157151>
- Pignatti, S. (1982). *Flora d'Italia*, Edagricole.
- POWO (2023). Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. <http://www.plantsoftheworldonline.org/> Retrieved 15 April 2023.
- Pulević, V. (2005). Material for vascular flora of Montenegro: supplement to "Conspectus florae Montenegrinae" (J. Rohlena). Republic Institute for Nature Protection (in Montenegrin).
- Rohlena, J. (1942). Conspectus florae Montenegrinae. *Preslia*, 20–21, 1–506.
- Romanov, R.E., Dragičević, S., & Troia, A. (2024). *Isoetes gymnocarpa* and *Utricularia* × *neglecta* – new taxa for Montenegro. *Acta Botanica Croatica*, 83(2). <https://doi.org/10.37427/botcro-2024-011>
- Salisbury, E. J. (1974). The reproduction of *Juncus tenuis* (*Juncus macer*) and its dispersal. *Transactions of the Botanical Society of Edinburgh*, 42(2), 187–190.
- Stešević, D., Luković, M., Caković, D., Ružić, N., Bubanja, N., & Šilc, U. (2017). Alien species in sand dune plant communities on Velika plaža in Ulcinj (Montenegro). *Periodicum biologorum*, 119(4), 239–249. <https://doi.org/10.18054/pb.v119i4.4917>
- Stešević, D., Kuzmić, F., Milanović, Đ., Stanišić-Vujačić, M., & Šilc, U. (2020). Coastal sand dune vegetation of Velika plaža (Montenegro). *Acta Botanica Croatica*, 79(1), 43–54. <https://doi.org/10.37427/botcro-2020-003>
- Stešević, D., Milanović, Đ., Stanišić-Vujačić, M., & Šilc, U. (2021). *Aristida oligantha* – a new alien species on the eastern Adriatic coast. *Acta Botanica Croatica*, 80(2), 217–220. <https://doi.org/10.37427/botcro-2021-019>
- Stroh, P. A., Leach, S. J., August, T. A., Walker, K. J., Pearman, D. A., Rumsey, F. J., Harrower, C. A., Fay, M. F., Martin, J. P., Pankhurst, T., Preston, C. D., & Taylor, I. (2014). A Vascular Plant Red List for England. *BSBI News*, 9, 1–193.
- Šilc, U., Mullaj, A., Alegro, A., Ibraliu, A., Dajić Stevanović, Z., Luković, M., & Stešević, D. (2016). Sand dune vegetation along the eastern Adriatic coast. *Phytocoenologia*, 46(4), 339–355.
- Šilc, U., Stešević, D., Rozman, A., Caković, D., & Kuzmić, F. (2019). Alien species and their impact on sand dunes in NE Adriatic. In: C.W. Finkl, & C. Makowski (Eds.), *Impacts of Invasive Species on Coastal Environments: Coasts in Crisis* (pp. 113–143). Springer, Cham.
- Šilc, U., Stešević, D., Luković, M., & Caković, D. (2020). Changes of a sand dune system and vegetation between 1950 and 2015 on Velika plaža (Montenegro, E Mediterranean). *Regional Studies in Marine Science*, 35, 101139. <https://doi.org/10.1016/j.rsma.2020.101139>
- Thiers, B. M. (updated continuously). Index Herbariorum. <https://sweetgum.nybg.org/science/ih/>, accessed September 2023.
- Tomaselli, V, Beccarisi, L, Cambria, S, Forte, L, Minissale, P, Sciadrello, S, Veronico, G, & Brullo, S. (2022). Validation of associations for the temporary ponds of the class *Isoeto-Nanojuncetea* in Puglia (southern Italy). *Mediterranean Botany*, 43, e80627. <https://doi.org/10.5209/mbot.80627>
- Troia, A., & Greuter, W. (2014). A critical conspectus of Italian *Isoetes* (Isoetaceae). *Plant Biosystems*, 148(1), 13–20. <https://doi.org/10.1080/11263504.2013.878409>
- Troia, A., & Greuter, W. (2015). A conspectus of and key to Greek *Isoetes* (Isoetaceae), based on a reassessment of Haussknecht's gatherings of 1885. *Willdenowia*, 45(3), 391–403.
- Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M., & Webb, D. A. (Eds.) (1972–1980). *Flora Europaea* (Vol. 3–5). Cambridge University Press.
- Verloove F. (2010). *Juncus dichotomus* (Juncaceae) in northwestern Italy, a xenophyte new to Europe. *Willdenowia*, 40(2), 173–178. <https://doi.org/10.3372/wi.40.40202>

## Appendix

Geographical coordinates (WGS84) and dates of species records: Species, longitude, latitude, date, habitat type, voucher number (TGU, BALU), and/or reléve number, observer/s and/or collector/s, (DS – Danijela Stešević, MSV – Milica Stanišić-Vujačić, ĐM – Đorđije Milanović):

***Isoetes histrix* s. l.:** 41.89434, 19.32877, 03.06.2021, Mediterranean temporary pond, TGU 2032985, Rel. No. 3621-1, DS & MSV; 41.89332, 19.33182, 03.06.2021, Mediterranean temporary pond, TGU 2032960, DS & MSV; 41.89345, 19.332, 03.06.2021, Mediterranean temporary pond, Rel. No. 3621-3, DS & MSV; 41.89608, 19.31878, 21.05.2021, Wet meadow, Rel. No. 21521-2, DS & MSV; 41.88996, 19.31995, 08.06.2021, Mediterranean temporary pond, TGU 2032961, Rel. No. 8621-5, DS & MSV; 41.90386, 19.28094, 19.05.2021, Wet meadow, Rel. No. 19521-3, DS & MSV; 41.90742, 19.26967, 08.06.2021, Mediterranean temporary pond, TGU 2032962, DS & MSV; 41.89387, 19.31432, 02.06.2021, Mediterranean temporary pond, observation, DS & MSV; 41.90865, 19.25963, 08.06.2021, Mediterranean temporary pond, observation, DS & MSV; 41.90945, 19.2568, 08.06.2021, Mediterranean temporary pond, TGU 2032963, DS & MSV; 41.894346, 19.328748, 03.06.2021, Mediterranean temporary pond, BALU DjM 02/01-002, Rel. No. 21Gj017, ĐM; 41.890293, 19.318302, 03.06.2021, Mediterranean temporary pond, BALU DjM 02/01-004, Rel. No. 21Gj021, ĐM; 41.879859, 19.364680, 14.05.2021, Wet meadow, BALU DjM 02/01-001, Rel. No. 21Gj008, ĐM

***Solenopsis laurentia* ssp. *gasparrinii*:** 41.89434, 19.32877, 03.06.2021, Mediterranean temporary pond, TGU 2032956, Rel. No. 3621-1, DS & MSV; 41.89332, 19.33182, 03.06.2021, Mediterranean temporary pond, Rel. No. 3621-2, DS & MSV; 41.89345, 19.332, 03.06.2021, Mediterranean temporary pond, Rel. No. 3621-3, DS & MSV; 41.8917, 19.33505, 03.06.2021, Mediterranean temporary pond, TGU 2032957, Rel. No. 3621-4, DS & MSV; 41.88909, 19.32218, 08.06.2021, Mediterranean temporary pond, Rel. No. 8621-3, DS & MSV; 41.894346, 19.328748, 03.06.2021, Mediterranean temporary pond, BALU DjM 178/10-001, Rel. No. 21Gj017, ĐM.

***Medicago doliata*:** 41.89927, 19.29064, 21.06.2021, Dune grassland, TGU 2032981, DS & MSV; 41.884, 19.35097, 21.05.2021, Dune grassland, TGU 2032983, DS & MSV;

***Myosotis laxa* ssp. *cespitosa*:** 41.88624, 19.33149, 15.05.2021, Dune grassland, TGU 2032984, DS & MSV; 41.89399, 19.33384, 21.05.2021, Dune grassland, TGU 2032985, DS & MSV; 41.88909, 19.32218, 08.06.2021, Mediterranean temporary pond, Rel. No. 8621-3, DS & MSV; 41.991802, 19.287593, 10.05.2022, BALU DjM 159/25-045, ĐM

***Avellinia festuroides*:** 41.89434, 19.32877, 03.06.2021, Mediterranean temporary pond, TGU 2032964, Rel. No. 3621-1, DS & MSV; 41.89332, 19.33182, 03.06.2021, wet meadow, Rel. No. 3621-2, DS & MSV; 41.89096, 19.31596,

08.06.2021, dune grassland, Rel. No. 8621-7, DS & MSV; 41.88996, 19.31995, 08.06.2021, Mediterranean temporary pond, Rel. No. 8621-5, DS & MSV; 41.90203, 19.28106, 17.06.2021, dune grassland, Rel. No. 17621-3, DS & MSV; 41.90607, 19.26658, 17.06.2021, dune grassland, TGU 2032965, DS & MSV; 41.90355, 19.27532, 17.06.2021, dune grassland, Rel. No. 17621-2, DS & MSV; 41.876881, 19.358625, 03.05.2021, Annual grassland, BALU DjM 66/17-001, Rel. No. 12Gj048, ĐM; 41.892648, 19.333243, 13.05.2021, Annual grassland, BALU DjM 66/17-002, Rel. No. 21Gj006, ĐM; 41.894346, 19.328748, 03.06.2021, Mediterranean temporary pond, BALU DjM 66/17-003, Rel. No. 21Gj017, ĐM; 41.887495, 19.344397, 14.05.2021, Annual grassland, BALU DjM 66/17-004, Rel. No. 21Gj007, ĐM; 41.898782, 19.279934, 14.05.2021, edge of Pine plantation on dunes, observer ĐM.

***Corynephorus divaricatus*:** 41.89757, 19.30147, 02.06.2021, dune grassland, TGU 2032975, DS & MSV; 41.89511, 19.30467, 02.06.2021, dune grassland, TGU 2032967, DS & MSV; 41.8864, 19.34561, 08.06.2021, dune grassland, Rel. No. 8621-1, DS & MSV; 41.87673, 19.36472, 08.06.2021, dune grassland, TGU 2032968, DS & MSV; 41.8876, 19.34259, 21.06.2021, dune grassland, TGU 2032970, DS & MSV; 41.9026, 19.27674, 21.06.2021, dune grassland, TGU 2032971, DS & MSV; 41.89387, 19.31432, 02.06.2021, dune grassland, observers, DS & MSV; 41.89784, 19.29799, 02.06.2021, dune grassland, Rel. No. 2621-2, DS & MSV; 41.89621, 19.30191, 02.06.2021, dune grassland, Rel. No. 2621-1, DS & MSV; 41.88624, 19.33149, 08.06.2021, dune grassland, observers, DS & MSV; 41.88875, 19.32254, 21.06.2021, dune grassland, observers, DS & MSV; 41.89927, 19.29064, 08.06.2021, dune grassland, observers, DS & MSV; 41.90063, 19.28619, 21.06.2021, dune grassland, TGU 2032973, DS & MSV; 41.87641, 19.36443, 29.06.2021, dune grassland, observers, DS & MSV; 41.88806, 19.34213, 08.06.2021, dune grassland, observers, DS & MSV; 41.89096, 19.31596, 08.06.2021, dune grassland, Rel. No. 8621-7, DS & MSV; 41.88965, 19.31972, 08.06.2021, dune grassland, Rel. No. 8621-4, DS & MSV; 41.89826, 19.29583, 08.06.2021, dune grassland, observers, DS & MSV; 41.89276, 19.31154, 08.06.2021, dune grassland, Rel. No. 8621-8, DS & MSV; 41.89525, 19.30431, 08.06.2021, dune grassland, TGU 2032967, Rel. No. 8621-9, DS & MSV; 41.89613, 19.30196, 02.06.2021, dune grassland, observers, DS & MSV; 41.89777, 19.29821, 02.06.2021, dune grassland, observers, DS & MSV; 41.887495, 19.344397, 14.05.2021, Annual grassland, BALU DjM 66/40-001, Rel. No. 21Gj007, ĐM; 41.900496, 19.284250, 12.05.2023, dune grassland, BALU DjM 66/40-002, Rel. No. 23Gj010, ĐM.

***Juncus tenuis*:** 41.87553, 19.36044, 17.06.2021, Wet place, TGU 2032978, DS & MSV

***Setaria parviflora*:** 41.91447, 19.24997, 15.07.2021, Ruderalised dry grassland, TGU 2032980, DS & MSV; 41.907156, 19.283072, 06.10.2022, Ruderalised dry grassland, BALU DjM 66/116-017, ĐM