

On a new *Molinia caerulea* grassland association in the central Apennines (Italy)

Giampiero Ciaschetti¹ , Giovanni Sburlino², Roberto Venanzoni¹ 

Key words: Purple Moorgrass, meadows, syntaxonomy, Abruzzo, karst plateau

Ključne besede: Modra stožka, travišča, sintaksonomija, Abruci, kraška planota

Abstract

Molinia caerulea meadows represent a well-known Natura 2000 Habitat with a large potential distribution area in Europe. As for Italy, at present they were recognized only in its Northern and Southern part. The present study analyzes for the first time a new *Molinia caerulea* plant community found in a karst montane plateau of Central Italy. Its floristic-sociological characteristics allow the authors to propose the new association *Centaureo jaceae-Molinietum caeruleae*, with two subassociation (*-typicum* and *-caricetosum hostianae*), along with one variant. Despite of these Purple moorgrass meadows show some affinities with some Western Balkan plant associations belonging to *Trifolio-Hordeetalia*, the study suggests that the new association should be better referred to the *Molinetalia caeruleae* order.

Izveček

Travniki z vrsto *Molinia caerulea* predstavljajo dobro znan Natura 2000 habitatni tip z veliko potencialno razširjenostjo v Evropi, vendar so v Italiji do sedaj znani le v severnem in južnem delu. V članku prvič analiziramo novo rastlinsko združbo z vrsto *Molinia caerulea*, ki uspeva na kraški planoti v osrednji Italiji. Njene floristično-sociološke lastnosti so avtorjem omogočile opis nove asociacije *Centaureo jaceae-Molinietum caeruleae* z dvema subasociacijama (*-typicum* in *-caricetosum hostianae*) z eno varianto. Kljub temu, da ti travniki modre stožke kažejo določeno podobnost z nekaterimi asociacijami z zahodnega Balkana, ki jih uvrščamo v red *Trifolio-Hordeetalia*, rezultati kažejo, da je novo asociacijo bolj smiselno uvrstiti v red *Molinetalia caeruleae*.

Corresponding author:
Giampiero Ciaschetti
E-mail: curls@libero.it

Received: 15. 3. 2025
Accepted: 9. 12. 2025



¹ Department of Chemistry, Biology and Biotechnologies, University of Perugia, Italy

² Ca' Foscari University of Venice, Italy

Introduction

Molinia caerulea (L.) Moench is a caespitose, perennial grass with a N-Eurasian distribution area (Oberdorfer, 2001; Aeschimann et al., 2004), also reported in some African countries and introduced into the USA and Canada (Dančák et al., 2012). It is widespread in open plant communities on moors, heaths, bogs, fens, wet meadows, cliffs and lake shores, always on at least seasonally wet or peaty ground (Tutin et al., 1980; Taylor et al., 2001).

It is present in Italy together with its congener *Molinia arundinacea* Schrank, previously known as *Molinia caerulea* (L.) Moench subsp. *arundinacea* (Schrank) K. Richt. in an earlier Italian flora (Conti et al., 2005). We refer here only to *Molinia caerulea*, known as *Molinia caerulea* (L.) Moench subsp. *caerulea* in the above-mentioned earlier flora.

The presence of *Molinia caerulea* in central Italy is documented in Lazio and Abruzzo (Bartolucci et al., 2024a), although with a punctiform distribution (Guarrera & Tammaro, 1996; Anzalone et al., 2010; Conti & Bartolucci, 2016; Lucchese, 2018; Conti et al., 2019).

This species develops plant communities which take its name, purple moorgrass meadows on mineral, damp and/or peaty soils. In Europe, these occur predominantly in temperate to subarctic zones and, in their typical aspect, correspond to the mown grasslands also known as litter meadows (Ellenberg, 2009; Mucina et al., 2016).

In Italy, *Molinia caerulea* communities are mainly present in the north, from the plains to the mountains (e.g. Poldini, 1973; Marchiori & Sburlino, 1982; Gerdol, 1987; Balátová-Tulačková & Venanzoni, 1989, 1990; Bracco et al., 1994, 1998; Bracco & Nola, 1995; Sburlino et al., 1995; Andreis & Zavagno, 1996; Minghetti & Pedrotti, 2000; Pedrotti, 2001, 2004; Riviaccio et al., 2021), while they are rare and highly localised along the peninsula to the south (Guido & Montanari, 1983; Barberis & Mariotti, 1981; Venanzoni, 1988; Bassi et al., 2015; Mariotti et al., 2015; Ciaschetti & Venanzoni, 2024). To date, there are no studies on *Molinia caerulea* communities in central Italy.

In terms of phytosociology, European purple moorgrass meadows are mostly attributed to the *Molinion caeruleae* alliance (*Molinietalia caeruleae*, *Molinio-Arrhenatheretea*) (e.g. Nowak & Fartmann, 2004; Havlová, 2006; Zelnik, 2011; Borhidi, et al. 2012; Mucina et al., 2016). Plant communities of this type have, however, also been attributed to other alliances belonging to the same class (Horvatić, 1963; Rivas-Martínez et al., 2001; Collaud et al., 2020) or to other classes (Oberdorfer, 1993; Kočí, 2010; Hájek & Hájková, 2011; Bergmeier, 2020).

Starting from Koch's fundamental work (1926) in

which he describes "*Molinietum caeruleae*", many associations have been proposed in Europe, for example Tüxen (1954), Philippi (1960), Passarge (1964), Marchiori & Sburlino (1982), Oberdorfer et al. (1967), Ellmauer & Mucina (1993), Redžić et al. (2013) etc. The syntaxonomy of these communities is, however, controversial. While some authors (Nowak & Fartmann, 2004; Havlová, 2006; Řezníčková, 2007; etc.) advocate a more comprehensive approach according to which, at least in large part and excluding acidophilous sphagnum fens, this association is still attributable to the association described by Koch (1926), others (Ellmauer & Mucina, 1993; Rivas-Martínez et al., 2001; Zelnik, 2011; Lafon et al., 2024) prefer instead an approach favouring differentiation into several *syntaxa* on an ecological and phytogeographic basis, similar to the earlier proposal by De Foucault & Géhu (1980).

The abovementioned association is, in fact, considered "central" to the *Molinion caeruleae* alliance (Oberdorfer, 1993) and it is therefore difficult to distinguish the diagnostic species of the many associations proposed (Sburlino et al., 1995).

In addition, in transitional situations between different phytogeographical areas as is often the case, the diagnostic species of a *syntaxon* are far from being exclusive to it. It is therefore often more appropriate to use geographical differentials to identify the different vegetation types (e.g. Werger & Van Gils, 1976; Poldini et al., 1998; Košir et al., 2008).

Excluding purple moorgrass meadows with sphagnum bogs associated with acidophilous fens (e.g. Brusa et al., 2006), the following *syntaxa* have been recognised in Italy as belonging to the *Molinion caeruleae* alliance:

- 1a. *Plantagini altissimae-Molinietum caeruleae* Marchiori et Sburlino 1982 (Marchiori & Sburlino, 1982; Sburlino et al., 1995);
- 1b. *Plantagini altissimae-Molinietum caeruleae* Marchiori et Sburlino 1982 *typicum* (Sburlino et al., 1995);
- 1c. *Plantagini altissimae-Molinietum caeruleae cladietosum marisci* Sburlino, Bracco, Buffa et Andreis 1995 (Sburlino et al., 1995);
- 2a. *Selino-Molinietum caeruleae* Kuhn 1937 (Gerdol, 1987; Balátová-Tulačková & Venanzoni, 1989; Sburlino et al., 1995);
- 2b. *Selino-Molinietum caeruleae inuletosum salicinae* Balátová-Tulačková et Venanzoni 1989 (Balátová-Tulačková & Venanzoni, 1989);
- 3a. *Succiso-Molinietum caeruleae* (Kovacs 1962) Soò 1969 (Minghetti & Pedrotti, 2000; Pedrotti, 2001);
- 3b. *Succiso-Molinietum caeruleae caricetosum elatae* Minghetti et Pedrotti 2000 (Minghetti & Pedrotti, 2000);

4. *Junco-Molinietum caeruleae* Preising 1951 (Barberis & Mariotti, 1981; Pedrotti, 2004; Venanzoni, 1988);
5. *Juncus acutiflorus* and *Molinia caerulea* community (Bracco & Nola, 1995; Sburlino et al., 1995). It was included in the *Juncion acutiflori* alliance, but according to Mucina et al. (2016) this alliance should be considered invalid according to Art. 2b of the ICPN (Theurillat et al. 2020) and should instead be attributed to *Molinion caeruleae*.
6. *Gentiano asclepiadeae-Molinietum caeruleae* Oberdorfer 1957 em. Oberdorfer et al. 1967 (Canullo et al., 1994; Vanacore Falco & Venanzoni, 2009);
7. *Stachyo-Molinietum* Passarge 1964 (Canullo et al., 1994, without data). This association was reported by Pedrotti (2003) too, but the author himself changed this attribution to *Junco-Molinietum* in a later paper (Pedrotti, 2004).
8. *Molinietum caeruleae* Koch 1926 (Montacchini, 1982);
9. *Trollio europaei-Molinietum coeruleae trifolietosum medii* de Foucault 1976 (Balátová-Tulačková & Venanzoni, 1990).

In the Dinaric Alps, *Molinia caerulea* meadows have been attributed both to the *Trifolio-Hordeetalia* order in areas with a submediterranean climate (Horvatić, 1963; Gaži-Baskova, 1963; Ritter-Studnička, 1972; Hulina, 2007) and to *Molinion caeruleae* in the montane belt, especially on siliceous substrates (e.g. Milanović et al., 2015). The *Molinietalia* order is associated with the temperate to subarctic zones of Europe, while the *Trifolio-Hordeetalia* order is limited to the Apennine and Balkan peninsulas (Mucina et al., 2016), although some authors also report it in a number of areas in Western Europe (e.g. Mercadal, 2020).

The floristic and vegetational affinities between the Apennines and the Balkan Peninsula have also been highlighted numerous times for other vegetation types (e.g. Biondi et al., 2002, 2006; 2014; Terzi et al., 2010; Di Pietro, 2011; Taffetani et al., 2012; Blasi & Del Vico, 2012; Terzi & D’Amico, 2016) and this affinity underlies the placement of both these geographical areas in the Apennine-Balkan phytogeographical province of the Euro-siberian region (Rivas-Martinez et al., 2004).

Molinia caerulea meadows are characterised by a high floristic diversity and rich fauna and are therefore of exceptional importance for the protection of species and biotopes. Moreover, their extent is now considerably reduced due to agricultural drainage and the use of fertilisers, as well as the abandonment of traditional mowing practices (Sburlino et al., 1995; Nowak & Fartmann, 2004, Nowak et al. 2015). The conservation of at least some complexes of the vast, unfertilised *Molinia caerulea*

grasslands is, in fact, one of the most pressing nature conservation problems in Europe (Ellenberg, 2009). In temperate transitional climates, namely those corresponding to the Submediterranean variant of the Temperate bioclimate according to the classification of Rivas-Martinez et al. (2011), also widespread in Abruzzo, *Molinia caerulea* communities are even more important in terms of conservation as they are home to numerous rare and endangered species at national or regional level (Venanzoni, 1988).

The *Molinia caerulea* grassland communities are mostly referable to the EU habitat “6410: *Molinia* meadows on calcareous, peaty or clayey-silt-laden soils (*Molinion caeruleae*)”, which is in poor condition in almost all biogeographical regions where it occurs (<https://www.biodiversity.europa.eu/habitats/6410> accessed on December 14th 2024).

This paper will attempt to answer the following questions:

1. do the *Molinia caerulea* meadows in the central Apennines have their own syntaxonomic autonomy, or are they related to any of the associations already found in Italy?
2. are the *Molinia caerulea* meadows in the central Apennines to be attributed to the *Molinion caeruleae* alliance or, like other meadows on temporarily flooded soils on central Italy’s karst plateaus, to the endemic *Ranunculum velutini* alliance of the *Trifolio-Hordeetalia* order (see Pedrotti & Sanesi, 1969; Pedrotti, 1976, 2019; Canullo et al., 1988; Pedrotti et al., 1992; Venanzoni, 1992; Pirone, 1997; Biondi et al., 1999; Tardella & Di Agostino, 2020)?

Methods

The vegetation studied was surveyed in the Sirente-Velino Regional Park, in the province of L’Aquila (Abruzzo), in an altitudinal range between 1300 and 1350 m a.s.l. Morphologically, this constitutes the central endorheic depression of the three making up the vast “Altopiano delle Rocche-Piano di Ovindoli” system of tectonic-karst plateaus. The karst nature of the area is evidenced by the presence of sinkholes through which the waters of the plain are drained, resulting in wide fluctuations in the water table throughout the year. The geological substrate consists of Plio-Quaternary sediments of fluvio-lacustrine origin. The soils are deep, with a scarce stony content, moderately fine texture and a reaction sub-acidic on the surface and neutral at depth.

Analysed over a thirty-year period, the bioclimate of the Rocca di Mezzo station (1329 m a.s.l.) is Oceanic Temperate, Submediterranean variant, with an Upper

Supra-Temperate thermo-type and a Lower Humid ombro-type, according to the bioclimatic classification of Rivas-Martínez et al. (2011). The Index of Continentality (IC = difference between the average temperature of the hottest month and the coldest month) is around 17°C, quite a high value compared to the rest of peninsular Italy (Pesaresi et al., 2017).



DISTRIBUTION OF THE MAIN *MOLINIA CAERULEA* PLANT COMMUNITIES IN ITALY

1. *Plantagini altissimae-Molinietum caeruleae*
2. *Selino-Molinietum caeruleae*
3. *Succiso-Molinietum caeruleae*
4. *Junco-Molinietum caeruleae*
5. *Juncus acutiflorus* and *Molinia caerulea* community
6. *Gentiano asclepiadeae-Molinietum caeruleae*
7. *Stachyo-Molinietum*
8. *Trollio europaei-Molinietum caeruleae*

Figure 1: Study area (*) and distribution of the main *Molinia caerulea* plant communities reported in Italy.

Slika 1: Raziskovano območje (*) in razširjenost glavnih rastlinskih združb z vrsto *Molinia caerulea* v Italiji.

In the summers of 2023 and 2024, 17 phytosociological relevés were carried out using the classical phytosociological method of the Zurich-Montpellier school (Braun-Blanquet, 1964). The 83 x 17 matrix obtained was transformed according to Van der Maarel (1979) and used for numerical processing employing the Syntax

2000 software (Podani, 2001). For the cluster analysis, the chord distance was used as the distance coefficient and the complete linkage as the clustering strategy. Type NMDS (Non Metric Multidimensional Scaling) ordination was used.

In order to assess whether Abruzzo's *Molinia caerulea* communities are closer to the associations/subassociations of northern Italy and the Alps, or to those described in the territories located further south, namely southern Italy and the western Balkans, a Minimum Spanning Tree was performed (again with the chord distance as the dissimilarity coefficient) on the synoptic table obtained with the data from the associations/subassociations found in these geographical areas (data sources in Appendix 1). With regard to the Balkans, the *Molinio-Lathyretum pannonicum* associations in Croatia and Bosnia (Horvatić, 1963; Gaži-Baskova, 1963; Ritter-Studnička, 1954, 1972) and the *Lathyro pannonicum-Molinietum caeruleae* association in Serbia (Tatić et al., 1988; Ačić et al., 2013) were considered, together with the related subassociations, with the exception of *Molinio-Lathyretum pannonicum klaseetosum lycopifoliae* as this refers to grasslands not dominated by *Molinia caerulea*.

Based on the results of the Minimum Spanning Tree and using the same methods as before, a further cluster analysis and NMDS sorting were carried out within the group of associations shown to be closest to those in Abruzzo.

A number of authors consider some of the associations to be synonyms, as in the case of *Selino carvifoliae-Molinietum caeruleae* Kuhn 1937, considered a synonym for *Carici davallianae-Molinietum caeruleae* Špáníková 1978 (www.geobot.org.ua/syntaxonomy/300/ accessed on December 12th 2024), and other authors propose considerably restricting the number of associations described (e.g. Havlová, 2006). We, however, deemed it appropriate to treat them separately, both because they represent geographical variations that could prove significant and because it is not the purpose of this paper to review the nomenclature of the communities considered.

Phytosociological tables consisting of a single relevé were not considered (e.g. Balátová-Tulačková & Venanzoni, 1990). For Bosnia, the data published in Ritter-Studnička (1972) were used, although the tables are not complete.

For the nomenclature of the species, the checklist of the Italian flora (Bartolucci et al., 2024a, 2024b; Galasso et al., 2024a, 2024b) was adopted. The syntaxonomic framework down to alliance level refers to the checklist of European vegetation (Mucina et al., 2016). As far as possible, the authors of the *syntaxa* names conform to Izzo (2002).

Figure 4) than with those in northern Italy and the Alps (numbers 11 to 24 in Figure 4). Although positioned close to the other “southern” communities in the diagram, the Calabrian phytocoenoses are linked to them with quite a high dissimilarity value. This is probably due to the particularly lithological characteristics of the Calabrian phytocoenoses, which grow on granitic-crystalline substrates.

The results of the MST make it possible to analyse the similarities/divergences between the communities in central-southern Italy and those on the Balkan Peninsula in greater detail.

The cluster analysis of the *Molinia caerulea* relevés in central-southern Italy and the western Balkans (Figure 5) shows two main groups (A and B in Figure 5 and 6), the second of which (B) consists of the relevés from Bosnia-Herzegovina. In the first cluster, the Calabrian relevés (A2) are shown to be independent of the rest of cluster A (A1), which in turn is divided into two sub-clusters: A1a, consisting of the Abruzzo and Serbia relevés, and A1b, including all the subassociations in the Croatian *Molinio-Lathyretum*.




Figure 5: Cluster analysis of the *Molinia caerulea* meadows in central-southern Italy and western Balkans: 1–2 = data from Abruzzo; 3 to 6 = *Molinio-Lathyretum pannonicum* from Croatia; 7–8 = *Molinio-Lathyretum pannonicum* from Bosnia; 9 = *Lathyro pannonicum-Molinietum caeruleae* from Serbia; 10 = *Junco effusi-Molinietum caeruleae* from Calabria (numbers as in Figure 4 and 6). Detailed data source in Appendix 1.

Figure 5: Klastrska analiza travnikov z vrsto *Molinia caerulea* v osrednji-južni Italiji in na zahodnem Balkanu: 1–2 = podatki iz Abrucov; 3 do 6 = *Molinio-Lathyretum pannonicum* s Hrvaške; 7–8 = *Molinio-Lathyretum pannonicum* iz Bosne; 9 = *Lathyro pannonicum-Molinietum caeruleae* iz Srbije; 10 = *Junco effusi-Molinietum caeruleae* iz Kalabrije (številke so kot v slikah 4 in 6). Natančni podatki o viru podatkov so predstavljeni v prilogi 1.

The NMDS ordination (Figure 6) confirms a separation of both clusters A and B and subclusters A1 and A2 along the first axis. The second axis not only emphasises the opposite positioning of A2 and B in two-dimensional space, it also indicates the sequential placement of subclusters A1a and A1b and, within the former, A1a1 and A1a2.




Figure 6: NMDS ordination of the *Molinia caerulea* meadows in central-southern Italy and western Balkans: 1–2 = data from Abruzzo; 3 to 6 = *Molinio-Lathyretum pannonicum* from Croatia; 7–8 = *Molinio-Lathyretum pannonicum* from Bosnia; 9 = *Lathyro pannonicum-Molinietum caeruleae* from Serbia; 10 = *Junco effusi-Molinietum caeruleae* from Calabria (numbers as in Figure 4–5). Detailed data source in Appendix 1.

Figure 6: NMDS ordinacija travnikov z vrsto *Molinia caerulea* v osrednji-južni Italiji in na zahodnem Balkanu: 1–2 = podatki iz Abrucov; 3 do 6 = *Molinio-Lathyretum pannonicum* s Hrvaške; 7–8 = *Molinio-Lathyretum pannonicum* iz Bosne; 9 = *Lathyro pannonicum-Molinietum caeruleae* iz Srbije; 10 = *Junco effusi-Molinietum caeruleae* iz Kalabrije (številke so kot v slikah 4 in 6). Natančni podatki o viru podatkov so predstavljeni v prilogi 1.

Discussion

In the southern and south-eastern European context, the *Molinia caerulea* phytocoenoses of Abruzzo (nos. 1 and 2 in Figure 4) are markedly different from those in northern Italy and the Alps (nos. 11–24 in Figure 4), although they share some *Molinion* and *Molinietalia caeruleae* diagnostic species, such as *Succisa pratensis*, *Serratula tinctoria* and *Sanguisorba officinalis*. It is, however, important to note that these species are also present, with varying frequencies, in the Balkan associations (nos. 3–9 in Figure 4).

The affinity between the Abruzzo and Balkan relevés is evidenced by the common presence of several species that are absent or poorly represented in the other phytocoenoses considered (see Appendix 1), such as *Lathyrus pannonicus* s.l., *Cynosurus cristatus*, *Trifolium repens* and *Trifolium montanum* s.l. Restricting the comparison to the Abruzzo and Balkan communities alone, the former share some species such as *Linum catharticum*, *Potentilla erecta* and *Filipendula vulgaris* only with the Serbian *Lathyro-Molinietum caeruleae*, and other species such as *Sesleria uliginosa*, *Carex hostiana* and *Festuca rubra* s.l. only with *Molinio-Lathyretum pannonicum* from Croatia and Bosnia-Herzegovina.

Looking at the cluster analysis and NMDS ordination of the “southern” *Molinia* grasslands investigated in the present study (Figs. 5–6), it can be seen that cluster B, relating to the Bosnia-Herzegovina formations is isolated

at the top of the graph, probably due to the fact that incomplete synoptic columns were used (16 and 24 species, respectively). A considerable distance also separates the communities in Calabria (A2 in Figure 5–6) from the others. This is due to the different nature of the substrates, which, in the case of the Calabrian rocks, determine greater water retention and soils with an acidic pH (Venanzoni, 1988).

With regard to cluster A1, the Abruzzo phytocoenoses (A1a1 in Figs. 5–6) is more closely linked to the Serbian *Lathyro pannonici-Molinietum caeruleae* (A1a2) than to the Croatian *Molinio-Lathyretum pannonicum* (A1b).

Despite apparent similarities in nomenclature, the syntaxonomic placement of the two Balkan associations is very different. Like most of the European associations, the Serbian *Lathyro-Molinietum caeruleae* is included in the *Molinion caeruleae* (Ačić et al., 2013), while the *Molinio-Lathyretum pannonicum* in Croatia and Bosnia-Herzegovina is attributed to the *Molinio-Hordeion secalini* (*Trifolio-Hordeetalia* order) (Horvatić, 1963; Ritter-Studnička, 1972). This different syntaxonomic positioning reflects the bioclimatic diversity of the areas concerned: sub-coastal areas in the colline belt with a submediterranean climate for *Molinio-Lathyretum pannonicum*; montane areas with a more continental climate for *Lathyro-Molinietum caeruleae*. As confirmation of this, among the species considered diagnostic of *Molinio-Hordeion secalini* and *Trifolio-Hordeetalia* (Horvatić, 1963; Šilc et al., 2014), only *Carex distans*, a species with a broad ecology, is present in the latter association.

The affinity of the Abruzzo relevés with those of the Balkans is primarily due to the latitudinal gradient which unites the Italian and Balkan peninsulas. Moreover, the central Apennine plateaus are karst in nature, as are the areas of the Balkan from which the relevés considered originate. On the other hand, the distinctly montane and moderately continental climate of the surveyed area explains its greater similarity to the Pešter Plateau (Serbia), rather than to the subcoastal areas of Croatia and Bosnia.

In the relevés presented here, as with the Serbian *Lathyro pannonici-Molinietum caeruleae*, the diagnostic species of the *Trifolio-Hordeetalia* are almost absent, while, as mentioned above, those diagnostic of the *Molinion caeruleae* and *Molinietalia caeruleae* are quite well represented.

Finally, several species are present in the Abruzzo relevés but absent from those from the Balkans: *Genista tinctoria*, *Carex flacca*, *Galium verum*, *Plantago media*, *Vicia cracca* and *Ranunculus polyanthemoides*. On the contrary, other species are present in the Balkan relevés and absent in those from the central Apennines: *Cirsium canum*, *Trifolium patens*, *Lysimachia nummularia*, *Anacamptis palustris*, *Oenanthe fistulosa*, etc. This gives the Abruzzo phyto-

coenoses a certain degree of autonomy, as shown by the separation of subcluster A1a1 in Figs. 5–6.

As can be seen, the *Molinia caerulea* communities studied are differentiated in two main aspects. The first (cluster I in Figs. 2–3) is associated with the more mesophilous situations, in contact with surrounding hay meadows growing on soils with a lower water content. The species differentiating this cluster are, in fact, typical of little to moderately fertilised hay meadows and/or pastures (e.g. *Genista tinctoria*, *Filipendula vulgaris*, *Plantago media* subsp. *media*, *Lathyrus pannonicus* subsp. *asphodeloides*, *Poa sylvicola*, *Trifolium montanum* subsp. *rupestre*). The meso-xerophilous species are even more prominent in subcluster Ib, which also includes species such as *Klasea lycopifolia*, *Thesium humifusum*, *Bromopsis erecta*, *Brachypodium genuense* and *Euphorbia gasparrini* subsp. *samnitica*. Of these, the last two species are endemic to Italy, while *Klasea lycopifolia* is a rare species protected by the European Habitats Directive 92/43 EEC and the subject of specific conservation actions in the study area (Gigante et al., 2014; Di Martino et al., 2016).

Cluster II (Figure 2–3), on the other hand, relates to edaphic situations with a shallower water table and higher content of organic matter, bordering on the low fen vegetation of the *Caricetalia davallianae* Br.-Bl. 1949 order described in Ciaschetti et al. (2024). The cluster is differentiated by a number of peatland or otherwise hygrophilous species: *Carex hostiana*, *Juncus inflexus* subsp. *inflexus*, *Dactylorhiza incarnata* subsp. *incarnata*, *Juncus articulatus* subsp. *articulatus*, *Carex davalliana*, *Carex nigra* subsp. *nigra*. In other areas, *Carex hostiana* in particular (also the most frequent) has already been used to distinguish autonomous associations, subassociations or their variants, together with *Carex davalliana* and other alkaline fen-related species (e.g. Koch, 1926; Klötzli, 1969; Havlová, 2006; Borhidi et al., 2012).

In light of these considerations, we consider it appropriate to classify the phytocoenoses studied into *Molinion*, *Molinietalia*, *Molinio-Arrhenatheretea* and, on the basis of their floristic-sociological autonomy, we propose the following new association:

CENTAUREO JACEAE-MOLINIETUM CAERULEAE Ciaschetti, Sbrulino et Venanzoni *ass. nova hoc loco* (holotypus rel. 1 of Table 1 this paper).

Description: this association refers to the *Molinia caerulea* dominated meadows found in flat areas of the karst montane plateaus of the central Apennines, on fine-textured, deep, temporary flooded soils, sub-acid on the surface and neutral in depth, generated on fluvio-lacustrine deposits.

Structure and floristic composition: it takes the form of a continuous hay meadow whose physiognomy is determined by the clearly dominant *Molinia caerulea*, accompanied by, among the most abundant species, *Succisa pratensis*, *Sesleria uliginosa*, *Serratula tinctoria*, *Festuca rubra* subsp. *commutata*, *Carex flacca* subsp. *flacca* and *Potentilla erecta*. *Klasea lycopifolia* is also very abundant locally. In particular, in the karst highlands of the central Apennines, *Sesleria uliginosa*, a species associated with moist soils (Sburlino et al., 1995; Foggi et al., 2001; Aeschmann et al., 2004) but with “a wide ecological amplitude with respect to moisture gradient” (Gonda & Dítě, 2011), had already been indicated as diagnostic of the meso-xerophilous *Cirsio acaulis*-*Seslerietum uliginosae* pasture association, (Biondi et al., 1992; Ciaschetti et al., 2006).

Diagnostic species: *Molinia caerulea*, *Centaurea jacea* subsp. *jacea*, *Rhinanthus minor*, *Ranunculus polyanthemoides*, *Sesleria uliginosa*. The last of these can be considered as a characteristic species at regional level (*sensu* Mucina, 1993).

Syntaxonomy: the association is included in the *Molinion caeruleae* alliance (*Molinietalia caeruleae*, *Molinio-Arrhenatheretea*). Several diagnostic species of these *syntaxa* are, in fact, very well represented. These include *Succisa pratensis*, *Sanguisorba officinalis* and *Serratula tinctoria*. The high edaphic humidity and biogeographical position in which these communities develop, namely the Alpine-Caucasian subregion of the Eurosiberian region (Rivas-Martinez et al., 2004), also justify this positioning.

Chorology: the association is so far known only for the Sirente-Velino Regional Park (Abruzzo, central Italy).

EU Reference Habitat: 6410 *Molinia* meadows on calcareous, peaty or clayey-silt-laden soils (*Molinion caeruleae*) (Blasi et al., 2010, available at <http://vnr.unipg.it/habitat>).

Conservation status, pressures and threats: the conservation status can be considered as relatively good. The area is mown for hay production once a year in the second half of July and no artificial fertilisation is used. After mowing, cattle grazing is permitted, but to such a modest extent that it does not pose an imminent threat.

CENTAUREO JACEAE-MOLINIETUM CAERULEAE TYPICUM Ciaschetti, Sburlino et Venanzoni *subass. nova hoc loco* (Table 1 rel. 1–9; *holotypus* rel. 1 of Table 1 this paper)

Description: this is the most widespread aspect in the study area, associated with the less hygrophilous aspects of the association, in contact with the meadows and pastures on areas not subject to flooding or only partially flooded. The lower water content enables greater mineralisation of

the organic matter in the soil, resulting in greater floristic richness, with an average of 34.6 species per relevé.

Structure and floristic composition: in addition to *Molinia caerulea*, the physiognomy is determined by *Succisa pratensis*, *Serratula tinctoria* and *Sesleria uliginosa*. The species of the *Molinion/Molinietalia* and *Molinio-Arrhenatheretea* are fairly well represented with an average of 6.2 and 9.7 entities per relevé respectively.

Diagnostic species: *Genista tinctoria*, *Filipendula vulgaris*, *Trifolium montanum* subsp. *rupestre*, *Lathyrus pannonicus* subsp. *asphodeloides*, *Plantago media* subsp. *media*, *Poa sylvicola*.

Klasea lycopifolia variant (Table 1 rel. 7–9)

Description: this variant has a richer presence of meso-xerophilous species from the little to moderately fertilised hay meadows and the pastures and can therefore be considered as an aspect of transition to the surrounding grassland communities in more elevated positions. Some similarities can be observed between this variant and *Lathyro asphodeloidis-Klaseetum lycopifoliae*, a type of meadow described in other localities close to the study area and attributed to the *Cynosurion cristati* alliance (Ciaschetti & Pirone, 2019). It should be noted that *Klasea lycopifolia* [= *Serratula lycopifolia* (Vill.) A.Kern] is also particularly abundant in *Molinio-Lathyretum pannonicum serratuletosum lycopifoliae*, the association's least hygrophilous subassociation, with an abundance of *Festuco-Brometea* species (Ritter-Studnička, 1972) which, as mentioned before, were not considered here as *Molinia caerulea* is absent.

Structure and floristic composition: In addition to the species determining the physiognomy of the association, *Klasea lycopifolia*, *Thesium humifusum*, *Brachypodium genuense* and *Bromopsis erecta* are also abundant.

CENTAUREO JACEAE-MOLINIETUM CAERULEAE CARICETOSUM HOSTIANAE Ciaschetti, Sburlino et Venanzoni *subass. nova hoc loco* (Table 1 rel. 10–17; *holotypus* rel. 14 of Table 1 this paper)

Description: This subassociation relates to the more hygrophilous aspects, in contact with the alkaline fens of the *Caricetalia davallianae*.

Structure and floristic composition: the physiognomy is established by *Molinia caerulea*, *Succisa pratensis*, *Gentiana pneumonanthe* subsp. *pneumonanthe*, *Centaurea jacea* subsp. *jacea*, *Carex flacca* subsp. *flacca*, *Potentilla erecta*, *Trifolium pratense* subsp. *pratense* and *Festuca rubra* subsp. *commutata* are also abundant.

Diagnostic species: *Carex hostiana* (regional characteristic *sensu* Mucina, (1993)). *Juncus inflexus* subsp. *inflexus*, *Dactylorhiza incarnata* subsp. *incarnata*, *Juncus articulatus*

subsp. *articulatus*, *Carex davalliana* and *Carex nigra* subsp. *nigra*. There is a certain depletion of diagnostic species in relevés 10–13, corresponding to subcluster IIa in Figure 2.

Overall, the vegetation studied can be classified as follows:

Syntaxonomic scheme

MOLINIO-ARRHENATHERETEA Tüxen 1937

MOLINIETALIA CAERULEAE Koch 1926

Molinion caeruleae Koch 1926

Centaureo jaceae-Molinietum caeruleae Ciaschetti, Sburlino et Venanzoni *ass. nova hoc loco*

– *typicum* Ciaschetti, Sburlino et Venanzoni *subass. nova*

– variant with *Klasea lycopifolia*

– *caricetosum hostianae* Ciaschetti, Sburlino et Venanzoni *subass. nova*

List of the syntaxa not quoted in the syntaxonomic scheme:

Carici davallianae-Molinietum caeruleae Špániková 1978

Cirsio acaulis-Seslerietum uliginosae Biondi, Ballelli, Allegranza, Frattaroli et Taffetani 1992 corr. Ciaschetti et al. 2006

Cynosurion cristati Tüxen 1947

Festuco-Brometea Br.-Bl. et Tüxen ex Soo 1947

Gentiano asclepiadeae-Molinietum caeruleae Oberdorfer 57 em. Oberdorfer et al. 1967

Juncion acutiflori Br.-Bl. 1947

Junco-Molinietum caeruleae Preising 1951

Junco effusi-Molinietum caeruleae Tüxen 1954

Lathyro-asphodeloidis-Klaseetum lycopifoliae Ciaschetti et Pirone 2019

Lathyro pannonici-Molinietum caeruleae Tatić et al. ex Ačić, Šilc, Lakušić, Vukojičić et Dajić Stevanović 2013

Molinietum caeruleae Koch 1926

Molinietum caeruleae caricetosum hostianae Koch 1926

Molinietum caeruleae caricetosum paniceae Koch 1926

Molinietum caeruleae caricetosum tomentosae Koch 1926

Molinio-Lathyretum pannonicum Horvatić 1963

Molinio-Lathyretum pannonicum typicum Horvatić 1963

Molinio-Lathyretum pannonicum caricetosum davallianae Gaži-Baskova 1963

Molinio-Lathyretum pannonicum caricetosum paniceae Ritter-Studnička 1972

Molinio-Lathyretum pannonicum serratuletosum lycopifoliae Ritter-Studnička 1972

Molinio-Lathyretum pannonicum salicetosum rosmarinifoliae Ritter-Studnička 1972

Molinio-Lathyretum pannonicum scilletosum pratensis Horvatić 1963

Molinio-Lathyretum pannonicum sieglingietosum Gaži-Baskova 1963

Plantagini altissimae-Molinietum caeruleae Marchiori et Sburlino 1982

Plantagini altissimae-Molinietum caeruleae typicum Sburlino, Bracco, Buffa et Andreis 1995

Plantagini altissimae-Molinietum caeruleae cladietosum marisci Sburlino, Bracco, Buffa et Andreis 1995

Ranunculion velutini Pedrotti 1978

Selino-Molinietum caeruleae Kuhn 1937

Selino-Molinietum caeruleae inuletosum salicinae Balátová et Venanzoni 1989

Stachyo-Molinietum Passarge 1964

Succiso-Molinietum caeruleae (Kovacs 1962) Soó 1969

Succiso-Molinietum caeruleae caricetosum elatae Minghetti et Pedrotti 2000

Trifolio-Hordeetalia Horvatić 1963

Trollio europaei-Molinietum caeruleae trifolietosum medii de Foucault 1976

Conclusions

The *Molinia caerulea* meadows described here are the first reported in central Italy, as are plant communities referable to the *Molinion caeruleae* alliance.

Molinia caerulea meadows in central Italy, together with those in Calabria, the southern Alps and the northern Apennines, represent important fragments of vegetation that can be considered relicts of the Ice Age, in common with other *Caricetalia davallianae* and *Magnocaricetalia* communities (e.g. Ciaschetti et al., 2021; Venanzoni et al., 2021; Ciaschetti et al., 2024).

The study shows how, in the transition zone between the submediterranean and temperate climates, high level of soil humidity and high-water table during the year can determine the establishment of phytocoenoses traced back to central European models, rather than to submediterranean communities of the *Trifolio-Hordeetalia* order. It is highly probable that this is also due to the altitudinal factor and degree of continentality, as is also the case in the Western Balkans.

From a strictly phytosociological point of view, the *Molinion caeruleae* alliance enriches the list of alliances to which the grasslands on wet and/or temporarily flooded soils of the central Apennine karst plateau can be attributed: *Salvio pratensis-Dactylidion glomeratae* Ubaldi et al. in Ubaldi 2003 (Venanzoni, 1992; Allegranza & Biondi, 2011; Biondi et al., 2011); *Cynosurion cristati* (Cortini Pedrotti et al., 1973; Pedrotti et al., 1992; Venanzoni, 1992; Pirone, 1997; Blasi et al., 2012; Ciaschetti & Pirone, 2019; Ciaschetti et al., 2021; *Ranunculion velutini* (Pedrotti, 1976, 2019; Pedrotti & Sanesi, 1969;


Canullo et al., 1988; Pedrotti et al., 1992; Venanzoni, 1992; Pirone, 1997; Biondi et al., 1999; Tardella & Di Agostino, 2020); *Potentillion anserinae* (Cortini Pedrotti et al., 1973; Pedrotti et al., 1992; Venanzoni, 1992; Pedrotti, 2019; Ciaschetti et al., 2021; Tardella & Di Agostino, 2020); *Mentho-Juncion inflexi* (Pedrotti, 1976; Pirone, 1997; Tardella & Di Agostino, 2020).

Research data availability

The data used in this study are available from the corresponding author upon reasonable request.

ORCID iDs

Giampiero Ciaschetti  <https://orcid.org/0000-0003-3314-8362>

Roberto Venanzoni  <https://orcid.org/0000-0002-7768-0468>

References

Ačić, S., Šilc, U., Lakušić, D., Vukojičić, S., & Dajić Stevanović, Z. (2013). Typification and correction of syntaxa from the class *Molinio-Arrhenatheretea* Tx. 1937 in Serbia. *Hacquetia* 12(2), 39–54. <https://doi.org/10.2478/HACQ-2013-0013>

Aeschimann, D., Lauber, K., Moser, D. M., & Theurillat, J. P. (2004). Flora alpina. Zanichelli.

Allegrezza, M., & Biondi, E. (2011). Syntaxonomic revision of the *Arrhenatherum elatius* grasslands of central Italy. *Fitosociologia*, 48(1), 23–40.

Andreis, C., & Zavagno, F. (1996). La vegetazione del Lago di Ganna, con particolare riferimento ai rapporti spaziali tra le cenosi dei *Molinietalia* e degli *Scheuchzerietalia palustris*. *Naturalista Valtellinese*, 7, 33–56.

Anzalone, B., Iberite, M., & Lattanzi, E. (2010). La Flora vascolare del Lazio. *Informatore Botanico Italiano*, 42(1), 187–317.

Balátová-Tulačková, E., & Venanzoni, R. (1989). Sumpf- und Feuchtrasengesellschaften in der Verlandungszone des Kalterer Sees (Lago di Caldaro), der Montiggler (Monticolo) Seen und in der Etsch (Adige) Aue, Oberitalien. *Folia Geobotanica et Phytotaxonomica*, 24, 253–295.

Balátová-Tulačková, E., & Venanzoni, R. (1990). Beitrag zur Kenntniss der Nass- und Feuchtwiesen in der montanen Stufe der Provinz Bozen (Bolzano), Italie. *Tuexenia*, 10, 153–171.

Barberis, G., & Mariotti, M., 1981. Ricerche geobotaniche sulle zone umide del Gruppo di Voltri (Appennino ligure-piemontese). *Archivio Botanico e Biogeografico Italiano*, 57(1-2), 50–91.

Bartolucci, F., Peruzzi, L., Galasso, G., Alessandrini, A., Ardenghi, N. M. G., Astuti, G., Bacchetta, G., Banfi, E., Barberis, G., Bernardo, L., Bouvet, D., Bovio, M., Calvia, G., Castello, M., Cecchi, L., Del Guacchio, E., Domina, G., Fascetti, S., Gallo, L., Gottschlich, G., Guarino, R., Gubellini, L., Hofmann, N., Iberite, M., Jiménez-Mejías, P., Longo, D., Marchetti, D., Martini, F., Masin, R. R., Medagli, P., Peccenini, S., Prosser, F., Roma-Marzio, F., Rosati, L., Santangelo, A., Scoppola, A., Selvaggi, A., Selvi, F., Soldano, A., Stinca, A., Wagen-sommer, R. P., Wilhalm, T., & Conti, F. (2024a). A second update to the checklist of the vascular flora native to Italy. *Plant Biosystems*, 158(2), 219–296. <https://doi.org/10.1080/11263504.2024.2320126>.

Bartolucci, F., Domina, G., Ballelli, S., Conti, F., Fortini, P., Del Guacchio, E., Di Iorio, E., Galasso, G., Gubellini, L., Hofmann, N., Laface, V. L. A., Lonati, M., Mazzacava, G., Nota, G., Pesaresi, S., Pinzani, L., Prosser, F., Quaranta, L., Selvi, F., Tiburtini, M., Tiburtini, R., Wilhalm, T., & Lastrucci, L. (2024b). Notulae to the Italian native vascular flora: 17. *Italian Botanist*, 17, 13–21. <https://doi.org/10.3897/italianbotanist.17.126693>.

Bassi, S., Bolpagni, S., Pezzi, G., & Pattuelli, M. (2015). Habitat di interesse comunitario in Emilia-Romagna. Regione Emilia-Romagna, Direzione Generale Ambiente, Difesa del Suolo e della Costa, Servizio Parchi e Risorse forestali.

Bergmeier, E. (2020). The vegetation of Germany – a cross-referenced conspectus of classes, orders and alliances based on the EuroVegChecklist. *Tuexenia*, 40, 19–32. <https://doi.org/10.14471/2020.40.024>.

Biondi, E., Allegrezza, M., Ballelli, S., Calandra, R., Crescente, M. F., Frattaroli, A. R., Gratani, L., Rossi, A., Taffetani, F. (1992). Indagini per una cartografia fitoecologica dell'altipiano di Campo Imperatore (Gran Sasso d'Italia). *Bollettino Associazione Italiana di Cartografia*, 86, 85–98.

Biondi, E., Ballelli, S., Allegrezza, M., Taffetani, F., Frattaroli, A. R., Guitian, J., & Zuccarello, V. (1999). La vegetazione di Campo Imperatore (Gran Sasso d'Italia). *Braun-Blanquetia*, 16, 53–115.

Biondi, E., Casavecchia, S., Pinzi, M., Allegrezza, M., & Baldoni, M. (2002). The syntaxonomy of the mesophilous woods of the central and Northern Apennines (Italy). *Fitosociologia*, 39(2): 71–94.

Biondi, E., Allegrezza, M., Casavecchia, S., Pesaresi, S. & Vagge I. (2006). Lineamenti vegetazionali e paesaggio vegetale dell'Appennino centrale e settentrionale. *Biogeographia*, 27, 35–129.

Biondi E., Allegrezza M., Mentoni M., 2011. Vegetational and geomorphological analyses of a small biotope particularly important for biodiversity in Central Apennine. *Fitosociologia*, 48(2), 109–122.

Biondi, E., Blasi, C., Allegrezza, M., Anzellotti, I., Azzella, M.M., Carli, E., Casavecchia, S., Copiz, R., Del Vico, E., Facioni, L., Galdenzi, D., Gasparri, R., Lasen, C., Pesaresi, S., Poldini, L., Sburlino, G., Taffetani, F., Vagge, I., Zitti, S., & Zivkovic, L. 2014. Plant communities of Italy: The vegetation Prodrome. *Plant Biosystems*, 148, 728–814. <https://doi.org/10.1080/11263504.2014.948527>.

Blasi, C., & Del Vico, E. (2012). High mountain vegetation of the Apennines. *Berichte der Reinhold-Tüxen-Gesellschaft*, 24, 179–194.

Blasi, C., Biondi, E., Copiz, R., Galdenzi, D., & Pesaresi, S. (Eds.). (2010). Manuale italiano di interpretazione degli habitat (Direttiva 92/43/CEE). Artiser.

Blasi, C., Tilia, A., Rosati, L., Del Vico, E., Copiz, R., Ciaschetti, G., & Burrascano, S. (2012). Geographical and ecological differentiation in Italian mesophilous pastures referred to the alliance *Cynosurion cristati* Tx. 1947. *Phytocoenologia*, 41(4), 217–229. <https://doi.org/10.1127/0340-269X/2011/0041-0504>.

Borhidi, A., Kevey, B., & Lendvai, G. (2012). Plant communities of Hungary. Akadémiai Kiadó.

Bracco, F., & Nola, P. (1995). La vegetazione acquatica e palustre dei lagoni di Mercurago. *Fitosociologia*, 29, 137–150.

Bracco, F., Buffa, G., & Sburlino, G. (1994). Confronto corologico tra le associazioni a *Molinia caerulea* (L.) Moench della Pianura Padana ed i sintipi Centro Europei: Primi Risultati. *Giornale Botanico Italiano*, 128(1), 464–464. <https://doi.org/10.1080/11263509409437268>.

Bracco, F., Buffa, G., Ghirelli, L., Sburlino, G., & Zuccarello, V. (1998). The phytosociological information and the management of

the upspring vegetation of river Sile Regional Park (Venetian Plain-Northern Italy). *Archivio Geobotanico*, 4(1), 51–57.

Braun-Blanquet, J. (1964). Pflanzensozioologie – Grundzüge der Vegetationkunde. Springer.

Brusa, G., Raimondi, B., & Cerabolini, B. (2006). La vegetazione della Riserva Naturale “Lago di Biandronno” (Lombardia, Italia settentrionale). *Fitosociologia*, 43(2), 111–128.

Canullo, R., Pedrotti, F., & Venanzoni, R. (1988). I prati umidi ed inondati dell’Alto Trigno (Molise, Italia). *Documents Phytosociologiques*, 11, 583–606.

Canullo, R., Pedrotti, F., & Venanzoni, R. (1994). La torbiera di Fivavé. In *Guida Escursione Società Italiana di Fitosociologia in Trentino (1-5 luglio 1994)*. Dipartimento di Botanica ed Ecologia Camerino (pp. 78–110).

Ciaschetti, G., & Pirone, G. (2019). *Lathyro asphodeloidis-Klaseetum lycopifoliae*, a new plant association in the alliance *Cynosurion cristati* Tüxen 1947 in Central Apennines. *Italian Botanist*, 7, 35–50. <https://doi.org/10.3897/italianbotanist.7.33361>.

Ciaschetti, G., & Venanzoni, R. (2024). #132. Annex I Habitat: 6410 *Molinia* meadows on calcareous, peaty or clayey-silt-laden soils (*Molinia caerulea*). In Rivieccio, G., Bagella, S., Bazan, G., Bonari, G., Brunco, V., Buffa, G., Candini, F., Cannucci, S., Caria, M. C., Ciaschetti, G., De Simone, L., Di Pietro, A., Fanfarillo, E., Ferrarato, M., Fiaschi, T., Gianguzzi, L., Lonati, M., Lorenzato, L., Marengo, G., Mainetti, A., Mascia, F., Mei, G., Nota, G., Iamonic, D., Patera, G., Perrino, E. V., Preo, S. M., Rocca, R., Stinca, A., Terranova, G., Troia, A., Venanzoni, R., & Angiolini, C. New national and regional Annex I Habitat records: from #123 to #138. *Plant Sociology*, 61(2), 73–84. <https://doi.org/10.3897/pls2024612/05>.

Ciaschetti, G., Pirone, G., Frattaroli, A. R. & Corbetta, F. (2006). La vegetazione del Piano di Pezza (Parco Naturale Regionale “Sirente-Velino”, Italia Centrale). *Fitosociologia*, 43(1), 67–84.

Ciaschetti, G., Pirone, G., & Venanzoni, R. (2021). Sedge vegetation of the ‘Major Highlands of Abruzzo’ (Central Italy): updated knowledge after new discoveries. *Plant Biosystems*, 155, 647–662. <https://doi.org/10.1080/11263504.2020.1801876>.

Ciaschetti, G., Paleskouskaya, S., & Venanzoni, R. (2024). Relicts of threatened biodiversity: the alkaline fens of the 7230 EU habitat on montane plateaus of Central Apennines (Italy). *Plants*, 13(10), 1282. <https://doi.org/10.3390/plants13101282>.

Collaud, R., Greffier, B., Ferrez, Y., & Bailly, G. (2020). Inventaire des végétations de Franche-Comté (d’après le Synopsis des groupements végétaux de Franche-Comté, Ferrez et al., 2011). Version avril 2020. Conservatoire botanique national de Franche-Comté – Observatoire régional des Invertébrés.

Conti, F., Abbate, G., Alessandrini, A., & Blasi, C. (2005). An annotated Checklist of the Italian Vascular Flora. Palombi.

Conti, F., & Bartolucci, F. (2016). The vascular flora of Gran Sasso and Monti della Laga National Park (Central Italy). *Phytotaxa*, 256, 1–119. <https://doi.org/10.11646/phytotaxa.256.1.1>.

Conti, F., Ciaschetti, G., Di Martino, L., & Bartolucci, F. (2019). An annotated checklist of the vascular flora of Majella National Park (Central Italy). *Phytotaxa*, 412, 1–90. <https://doi.org/10.11646/phytotaxa.412.1.1>.

Cortini Pedrotti, C., Orsomando, E., Pedrotti, F., & Sanesi, G. (1973). La vegetazione ed i suoli del Pian Grande di Castelluccio di Norcia (Appennino centrale). *Atti dell’Istituto Botanico e Laboratorio Crittografico Università di Pavia, Ser. 6(9)*, 155–249.

Dančák, M., Duchoslav, M., & Trávníček, B. (2012). Taxonomy and cytogeography of the *Molinia caerulea* complex in central Europe. *Preslia*, 84, 351–374.

De Foucault, B., & Géhu, J.-M. (1980). Essai synsystème et chorologique sur les prairies a *Molinia caerulea* et *Juncus acutiflorus* de l’Europe occidentale. *Colloques Phytosociologiques*, 7(1978), 135–164.

Di Martino, L., Conti, F., Bartolucci, F., Ciaschetti, G., & Di Cecco, M. (2016). Conservazione delle piante di interesse comunitario (Direttiva 92/43/CEE) dell’Appennino centrale: il progetto Liffa Floranet. *Notiziario della Società Botanica Italiana*, 0, 45–46.

Di Pietro, R. (2011). New Dry Grassland Associations from the Ausoni-Aurunci Mountains (Central Italy) – Syntaxonomical Updating and Discussion on the Higher Rank Syntaxa. *Hacquetia*, 10(2), 183–231. <https://doi.org/10.2478/v10028-011-0011-9>.

Ellenberg, H. 2009. Vegetation ecology in Central Europe, 4th Edition. Cambridge University Press.

Ellmayer, T., & Mucina, L. (1993). *Molinia-Arrhenatheretea*. In: L. Mucina, G. Grabherr & T. Ellmayer (Eds.). Die Pflanzengesellschaften Österreichs I. Anthropogene Vegetation. G. Fischer, (pp. 297–401).

Foggi, B., Nardi, E., Rossi, G. (2001). Nomenclatural notes and typification in *Sesleria* Scop. (Poaceae). *Taxon*, 50, 1101–1106. <https://doi.org/10.2307/1224726>.

Galasso, G., Conti, F., Peruzzi, L., Alessandrini, A., Ardenghi, N. M. G., Bacchetta, G., Banfi, E., Barberis, G., Bernardo, L., Bouvet, D., Bovio, M., Castello, M., Cecchi, L., Del Guacchio, E., Domina, G., Fascetti, S., Gallo, L., Guarino, R., Gubellini, L., Guiggi, A., Hofmann, N., Iberite, M., Jiménez Mejías, P., Longo, D., Marchetti, D., Martini, F., Masin, R. R., Medagli, P., Musarella, C. M., Peccenini, S., Podda, L., Prosser, F., Roma-Marzio, F., Rosati, L., Santangelo, A., Scoppola, A., Selvaggi, A., Selvi, F., Soldano, A., Stinca, A., Wagensommer, R. P., Wilhalm, T., & Bartolucci F. (2024a). A second update to the checklist of the vascular flora alien to Italy. *Plant Biosystems*, 158(2): 297–340. <https://doi.org/10.1080/11263504.2024.2320129>

Galasso, G., Domina, G., Bacchetta, G., Barberis, D., Bartolucci, F., Cancellieri, L., Ceschin, S., Ciaramella, D., Croce, A., Cuenca-Lombrana, A., Del Guacchio, E., Di Lernia, D., Fois, M., Fontana, D., Franzoni, J., Giacò, A., Laface, V. L. A., Lallai, A., Lonati, M., Lupoletti, J., Maccioni, A., Mascia, F., Mei, G., Morabito, A., Musarella, C. M., Pelella, E., Pica, A., Pinzani, L., Podda, L., Stinca, A., Varricchione, M., & Lastrucci, L. (2024b). Notulae to the Italian alien vascular flora: 17. *Italian Botanist*, 17, 43–53. <https://doi.org/10.3897/italianbotanist.17.126768>

Gaži-Baskova, V. (1963). Zajednica beskoljenke *Molinio-Lathyretum pannonicum* u Lici i Krbavi. *Agronomski Glasnik*, 25(3), 165–175.

Gerdol, R., 1987. Geobotanical investigations in the small lakes of Lombardy. *Atti dell’Istituto Botanico e Laboratorio Crittografico Università di Pavia, Ser. 7, vol. 6* (1987), 5–49.

Gigante, D., Alessandrini, A., Ballelli, S., Bartolucci, F., Conti, F., Ferri, V., Gubellini, L., Hofmann, N., Montagnani, C., Pinzi, M., Venanzoni, R., & Wagensommer, R. P. (2014). Schede per una Lista Rossa della Flora vascolare e crittogamica Italiana: *Klasea lycopifolia* (Vill.) Á.Löve et D.Löve. *Informatore Botanico Italiano*, 46(1), 93–152.

Gonda, M., & Dítě, D. (2011). The ecology and coenotic characteristics of *Sesleria uliginosa* in Slovakia and comparison with Czech Republic. *Zprávy České Botanické Společnosti, Praha*, 46, 161–192.

- Guarrera, P. M., & Tammara, F. (1996). La flora del M. Sirente e zone limitrofe (Appennino Abruzzese). *Annali di Botanica (Roma)*, 52, Suppl. 11, 267–381.
- Guido, M., & Montanari, C. (1983). Studio e cartografia della vegetazione cacuminale del Monte Aiona (Appennino ligure). *Archivio Botanico e Biogeografico Italiano*, 59(3/4), 105–131.
- Hájek, M., & Hájková, P. (2011). Vegetation of fens, transitional mires and bog hollows. In: M. Chytrý (Ed.), *Vegetace České republiky 3. Vodní a mokřadní vegetace. [Vegetation of the Czech Republic 3. Aquatic and wetland vegetation]* (pp. 614–704). Academia.
- Havlová, M. (2006). *Molinion* meadows in the Czech Republic. *Preslia*, 78, 87–101.
- Horvatić, S. (1930) Soziologische Einheiten der Niederungswiesen in Kroatien und Slavonien. *Acta Botanica Croatica*, 5(1), 57–118.
- Horvatić, S. (1963). Vegetacijska karta otoka Paga općim pregledom vegetacijskih jedinica hrvatskog primorja. *Prirodoslovna istraživanja serija Acta biologica*, 4(33), 5–181
- Hulina, N. (2007). Flora and vegetation of the Drežnica field. *Agronomski Glasnik*, 4, 255–276.
- Izco J. (2002). Authors of syntaxon names. *Itinera Geobotanica*, 15(1), 413–432.
- Klötzli, F. (1969). Die Grundwasserbeziehungen der Streu- und Moorwiesen im nördlichen Schweizer Mittelland. *Beitr. Geobot. Landesaufl.* 52, 1–296.
- Koch, W. (1926). Die Vegetationseinheiten der Linthebene unter Berücksichtigung der Verhältnisse in der Nordostschweiz. Systematisch-kritische Studie. *Jahrbuch der St Gallischen naturwissenschaftlichen Gesellschaft*, 61(2), 1–146.
- Kočí, M. (2010). Subalpine tall-forb and deciduous-shrub vegetation (*Mulgedio-Aconitetea*). In M. Chytrý (Ed.), *Vegetace České republiky. 1. Travinná a keříčková vegetace [Vegetation of the Czech Republic. 1. Grassland and Heathland Vegetation]* (pp. 91–131). Academia.
- Košir, P., Čarni, A., & Di Pietro, R. (2008). Classification and phytogeographical differentiation of broad-leaved ravine forests in southeastern Europe. *Journal of Vegetation Science*, 19, 331–342. <https://doi.org/10.3170/2008-8-18372>.
- Lafon, P., Argagnon, O., Belaud, A., Cateau, E., Causse, G., Corriol, G., Culat, A., Delassus, L., Dumoulin, J., Gaudillat, V., Goret, M., Mangeat, M., Millet, J., Noble, V., Paulin, D., Soucanye, & De Landevoisin, C.-A. (2024). Catalogue de la végétation de France métropolitaine (CatVeg). *Bulletin de la Société Botanique du Nord de la France, n° spécial*, 1–372.
- Lucchese, F. (2018). Atlante della Flora Vascolare del Lazio, cartografia, ecologia e biogeografia. Vol. 2. La flora di maggiore interesse conservazionistico. Regione Lazio, Direzione Capitale Naturale, Parchi e Aree Protette.
- Marchiori, S., & Sburlino, G. (1982). I prati umidi dell'anfiteatro morenico del Tagliamento (Friuli – Italia nord-orientale). *Documents Phytosociologiques*, 7, 199–222.
- Mariotti, M. G., Marsili, S., Pavarino, M., Poponessi, S., & Zotti M. (2015). Note sulle conoscenze botaniche della Riserva Naturale Orientata delle Agoraie e dei suoi dintorni (Appennino Ligure). *Bollettino dei Musei e degli Istituti biologici dell'Università di Genova*, 77, 24–57.
- Mercadal, G. (2020). Geobotanical characterization and phytosociological validation of various syntaxons of dall hygrophils prats (ord. *Deschampsietalia cespitosae*) and mesohygrophils (order *Trifolio-Hordeetalia*) from Catalonia and Western Europe. *Bulletti de la Institució Catalana d'Història Natural*, 84, 163–193. <https://doi.org/10.2436/20.1502.01.49>.
- Milanović, Đ., Brujić, J., Đug, S., Muratović, E., & Lukić Bilela, L. (2015). *Vodič kroz tipove staništa BiH (A guide to habitat types of BiH)*. Prospect C&S.
- Minghetti, P., & Pedrotti, F. 2000. La vegetazione del Laghetto delle Regole di Castelfondo (Trento). *Studi Trentini di Scienze Naturali, Acta biologica*, 74 (1997), 155–173.
- Montacchini, F. (1982). *Carta della vegetazione della Valle di Susa ed evidenziazione dell'influsso antropico. Scala 1:50.000*. C.N.R. Coll. Progr. Final. Promozione della qualità dell'ambiente. AQ/1/220.
- Mucina, L. (1993). Nomenklatorische und syntaxonomische Definitionen, Konzepte und Methoden. In: L. Mucina, G. Grabherr & T. Ellmauer (Eds.), *Die Pflanzengesellschaften Österreichs 1. Anthropogene Vegetation*. (pp. 19–28). G. Fischer.
- Mucina, L., Bultmann, H., Dierßen, K., Theurillat, J-P, Raus, T., Carni, A., Sumberova, K., Willner, W., Dengler, J., Garcia, R.G., et al. (2016). Vegetation of Europe: hierarchical floristic classification system of vascular plant bryophyte lichen and algal communities. *Applied Vegetation Science*, 19(S1), 3–264. <https://doi.org/10.1111/avsc.12257>.
- Nowak, B., & Fartmann, T. (2004). *Molinion caeruleae*. In M. Burkart et al. (Eds.), *Synopsis der Pflanzengesellschaften Deutschlands, Heft 9 Teil 2: Molinietalia* (pp. 62–82). Floristisch-soziologische Arbeitsgemeinschaft.
- Nowak T., Węgrzynek, B., & Tokarska-Guzik, B. (2015). Assets and threats to *Molinia* meadows (*Molinion caeruleae* alliance) on chosen natura 2000 areas in the eastern part of the silesian upland. *Acta Scientiarum Polonorum series Agricultura*, 14(4), 49–61.
- Oberdorfer, E. 1993. *Süddeutsche Pflanzengesellschaften, Band 3: Wirtschaftswiesen und Unkrautgesellschaften*. Gustav Fischer.
- Oberdorfer, E. 2001. *Pflanzensoziologische Exkursionsflora. 8th ed.* E. Ulmer.
- Oberdorfer, E., Görs, S., Korneck, D., Lohmeyer, W., Müller, T., Philippi, G., & Seibert, P. (1967). Systematische Übersicht der westdeutschen Phanerogamen- und Gefäßkryptogamen-Gesellschaften. *Schriftenreihe für Vegetationskunde*, 2, 6–62.
- Passarge, H. (1964). Pflanzengesellschaften der nordostdeutschen Flachlandes. *Pflanzensoziologie*, 13, 1–324.
- Pedrotti, F. (1976). Les prairies permanentes humides de l'Apennin central: phytosociologie et cartographie. *Colloques Phytosociologiques*, 5, 181–187.
- Pedrotti, F. (2001). Biotopkartierung im Trentino: Methoden und Resultate. *Sauteria*, 11, 61–74.
- Pedrotti, F. (2003). Vegetation changes in the Laghestel wetland basin (Trentino, Central Alps) from 1976 to 2001. Abstracts 46th Symp. I.A.V.S., 8-14/06/2003, Napoli, Italy: 171.
- Pedrotti, F. (2004). Ricerche geobotaniche al Laghestel di Piné. *Braun-Blanquetia*, 35, 1–55.
- Pedrotti, F. 2019. Flora e vegetazione della Palude di Colfiorito (Appennino Centrale, Italia). *Les Cahiers de Braun-Blanquetia*, 2, 1–118.

- Pedrotti, F., & Sanesi, G. (1969). Resoconto sulle escursioni dell'Appennino Umbro–Marchigiano (25–27 giugno 1968). *Mitteilungen der Ostalpin-Dinarischen pflanzensoziologischen Arbeitsgemeinschaft*, 9, 365–388.
- Pedrotti, F., Gafta, D., Manzi, A., & Canullo, R. (1992). Le associazioni vegetali della piana di Pescasseroli (Parco Nazionale d'Abruzzo). *Documents Phytosociologiques*, 14, 123–147.
- Pesaresi, S., Biondi, E., & Casavecchia, S. (2017). Bioclimates of Italy. *Journal of Maps*, 13(2), 955–960. <https://doi.org/10.1080/17445647.2017.1413017>.
- Philippi, G., (1960). Zur Gliederung der Pfeifengraswiesen im südlichen und mittleren Oberrheingebiet. *Beiträge zur naturkundlichen Forschung in Südwestdeutschland*, 19, 138–187.
- Pirone, G. 1997. *Il paesaggio vegetale di Rivisondoli, aspetti della flora e della vegetazione*. Edigrafital.
- Podani, J. 2001. *Syn-tax 2000. Computer program for data analysis in ecology and systematics*. Scientia Publishing.
- Poldini L., 1973. *Die Pflanzendecke der Kalkflachmoore in Friaul (Nordostitalien). Berichte des Geobotanischen Institutes der Eidg. Techn. Hochschule, Stiftung Rübel*, 51, 166–178.
- Poldini, L., Oriolo, G., & Mazzolini, G. (1998). The segetal vegetation of vineyards and crop fields in Friuli-Venezia Giulia (NE-Italy). *Studia Geobotanica*, 16, 5–32.
- Redžić, S., Trakić, S., & Barudanović, S. (2013). Patterns of vegetation diversity of grasslands and pastures – Crvanj Mt. (Herzegovina, Western Balkan). *Scientific Research and Essays*, 8(39), 1944–1965. <https://doi.org/10.5897/SRE11.1309>
- Řezníčková M. (2007). *Molinietum caeruleae* Koch 1926. – In: M. Chytrý (Ed.), *Vegetace České republiky. 1. Travinná a keříčková vegetace [Vegetation of the Czech Republic. 1. Grassland and Heathland Vegetation]* (pp. 210–213). Academia.
- Ritter-Studnička, H. (1954). Flora i vegetacija kraških polja Bosne i Hercegovine. *Godišnjak Bioloskog Instituta Sarajevo*, 1-2, 25–109.
- Ritter-Studnicka H. (1972). Neue Pflanzengesellschaften aus den Karstfeldern Bosniens und der Hercegovina. *Botanische Jahrbucher*, 92(1), 108–154.
- Rivas-Martínez, S., Fernández-González, F., Loidi, J., Lousá, M., & Penas, A. (2001). Syntaxonomical checklist of vascular plant communities of Spain and Portugal to association level. *Itinera Geobotanica*, 14, 1–341.
- Rivas-Martínez, S., Penas A., & Díaz T.E. (2004). *Biogeographic map of Europe*. Carthographic Service, University of León.
- Rivas-Martínez, S., Sáenz, S. R., & Penas A. (2011). Worldwide bioclimatic classification system. *Global Geobotany*, 1, 1–634
- Rivieccio, G., Aleffi, M., Angiolini, C., Bagella, S., Bazan, G., Bonini, F. et al. (2021). New national and regional Annex I Habitat records: from #26 to #36. *Plant Sociology*, 58(2), 77–98. <https://doi.org/10.3897/pls2021582/07>.
- Sburlino, G., Bracco, F., Buffa, G., & Andreis, C. (1995). I prati a *Molinia caerulea* (L.) Moench della Pianura padana: sintassonomia, sinecologia, sinecologia. *Fitosociologia*, 2, 67–87.
- Taffetani, F., Catorci, A., Ciaschetti, G., Cutini, M., Di Martino, L., Frattaroli, A.R., Paura, B., Pirone, G., Rismondo, M., & Zitti, S. (2012). The *Quercus cerris* woods of the alliance *Carpinion orientalis* Horvat 1958 in Italy. *Plant Biosystems*, 146(4), 918–953. <https://doi.org/10.1080/11263504.2012.682613>.
- Tardella, F.M., & Di Agostino, V. M. (2020). Vegetation of the “Altipiani di Colfiorito” wetlands (central Apennines, Italy). *Plant Sociology*, 57(2), 113–132. <https://doi.org/10.3897/pls2020572/04>.
- Tatić, B., Veljović, V., Petković, B., Stefanović, M., & Radotić, S. (1988). Ass. *Lathyreto-Molinietum coeruleae*-nova zajednica livadske vegetacije sa Pešterske visoravni-jugozapadna Srbija. *Glasnik Instituta za Botaniku i Botaničke Baste Univerziteta u Beogradu*, 12, 31–38.
- Taylor, K., Rowland, A. P., & Jones, H. E. (2001). *Molinia caerulea* (L.) Moench. Biological Flora of the British Isles NO. 216. *Journal of Ecology*, 89, 126–144.
- Terzi, M., Di Pietro, R., & D'Amico, F.S. (2010). Application of Indicator Species Analysis to communities with *Stipa austroitalica* Martinovsky and relevant syntaxonomic problems. *Fitosociologia*, 47(1), 3–29.
- Terzi, M., & D'Amico, F. S. (2016). Dry grasslands of *Hippocrepido glaucae-Stipion austroitalicae* in the Pollino Massif (Calabria, Italy). *Acta Botanica Croatica*, 75(1), 89–98. <https://doi.org/10.1515/botcro-2016-0013>.
- Theurillat, J. P., Willner, W., Fernández-González, F., Bültmann, H., Čarni, A., Gigante, D., et al. (2020). International Code of Phytosociological Nomenclature. 4th edition. *Applied Vegetation Science*, 24(1), e12491. <https://doi.org/10.1111/avsc.12491>.
- Tutin, T.G., Heywood, V.H., Burges, N.A., Moore, D.M., Valentine, D.H., Walters, S.M., & Webb, D.A. (Eds.). (1980). *Flora Europaea. Vol. 5*. Cambridge University Press.
- Tüxen, R. (1954). Pflanzengesellschaften und Grundwasserganglinien. *Angewandte Pflanzensoziologie*, 8, 64–98.
- Vanacore Falco, I., & Venanzoni, R. (2009). Indagine floristico-vegetazionale nel Sito d'Importanza Comunitaria “Talweg della Val Ferret” (IT 1204032) (Courmayeur, Aosta). *Revue Valdôtaine d'Histoire Naturelle*, 63, 23–62.
- Van Der Maarel, E. (1979). Transformation of cover-abundance values in phytosociology and its effects on community similarity. *Vegetatio*, 39(2), 97–114. <https://doi.org/10.1007/BF00052021>.
- Venanzoni, R. 1988. Contributo alla conoscenza dei prati umidi della Sila (Calabria-Italia). *Documents Phytosociologiques*, 11, 613–614.
- Venanzoni, R. (1992). I prati umidi e inondati dell'Alta Valle del Velino (Rieti-Italia Centrale). *Documents Phytosociologiques*, 14, 149–164.
- Venanzoni, R., Praleskouskaya, S., & Ciaschetti, G. (2021). Contribution to the Syntaxonomy of Rare Tall Sedge Community in Central Apennine (Umbria-Italy): I. *Caricetum buxbaumii*. *Flora Mediterranea*, 31 (Special Issue), 95–104. <https://doi.org/10.7320/FIMedit31SI.095>.
- Werger, M. J. A., & Van Gils, H. (1976). Phytosociological classification in chorological borderline areas. *Journal of Biogeography*, 3, 49–54. <https://doi.org/10.2307/3038098>.
- Zelnik, I. (2011). Wet meadows with Purple Moor-grass (*Molinia caerulea*) in Slovenia. *Acta Biologica Slovenica*, 54(2), 53–71. <https://doi.org/10.14720/abs.54.2.15479>.

Appendix 1

Data sources of Figures 4–6.

Data sources are in brackets.

- 1: *Centaureo jaceae-Molinietum caeruleae typicum* Ciaschetti, Sburlino et Venanzoni *subass. nova hoc loco* – Abruzzo, Italy;
- 2: *Centaureo jaceae-Molinietum caeruleae caricetosum hostianae* Ciaschetti, Sburlino et Venanzoni *subass. nova hoc loco* – Abruzzo, Italy;
- 3: *Molinio-Lathyretum pannonicum typicum* Horvatić 1963 – Croatia (Horvatić 1963);
- 4: *Molinio-Lathyretum pannonicum scilletosum pratensis* Horvatić 1963 – Croatia (Horvatić 1963);
- 5: *Molinio-Lathyretum pannonicum sieglingietosum* Gaži-Baskova 1963 – Croatia (Gaži-Baskova 1963);
- 6: *Molinio-Lathyretum pannonicum caricetosum davallianae* Gaži-Baskova 1963 – Croatia (Gaži-Baskova 1963);
- 7: *Molinio-Lathyretum pannonicum caricetosum paniceae* Ritter-Studnička 1972 – Bosnia (Ritter-Studnička 1972);
- 8: *Molinio-Lathyretum pannonicum salicetosum rosmarinifoliae* Ritter-Studnička 1972 – Bosnia (Ritter-Studnička 1972);
- 9: *Lathyro pannonicum-Molinietum caeruleae* Tatić et alii. ex Ačić, Šilc, Lakušić, Vukojičić, Dajić Stevanović 2013 – Serbia (Tatić et al. 1988);
- 10: *Junco effusi-Molinietum caeruleae* Tüxen 1954 – Calabria, Italy (Venanzoni 1988 sub *Junco-Molinietum caeruleae* Preising 1951);
- 11: *Junco effusi-Molinietum caeruleae* Tüxen 1954 – Liguria, Italy (Barberis & Mariotti 1981 sub *Junco-Molinietum caeruleae* Preising 1951);
- 12: *Juncus acutiflorus* and *Molinia caerulea* groupment – Piedmont, Italy (Bracco & Nola 1995);
- 13: *Gentiano asclepiadeae-Molinietum* Oberdorfer 57 em. Oberdorfer et alii 1967 – Trento Province, Italy (Cannullo et al. 1994);
- 14: *Succiso-Molinietum caeruleae* (Kovacs 1962) Soó 1969 – Trento Province, Italy (Minghetti & Pedrotti 2000);
- 15: *Selino-Molinietum caeruleae* Kuhn 1937 – Lombardy, Italy (Sburlino et al. 1995);
- 16: *Selino-Molinietum caeruleae inuletosum salicinae* Balatova et Venanzoni 1989 – South Tyrol, Italy (Balátová-Tulacková & Venanzoni 1989);
- 17: *Molinia caerulea* groupment – Lombardy, Italy (Andreis & Zavagno 1996);
- 18: *Plantagini altissimae-Molinietum caeruleae typicum* Sburlino, Bracco, Buffa & Andreis 1995 – Eastern Po Valley, Italy (Sburlino et al. 1995);

- 19: *Plantagini altissimae-Molinietum caeruleae cladietosum marisci* Sburlino, Bracco, Buffa & Andreis 1995 – Friuli, Italy (Sburlino et al. 1995);
- 20: *Trollio-Molinietum caeruleae* Guinochet 1955 – Jura Mountains, France/Switzerland (Oberdorfer 1993);
- 21: *Molinietum caeruleae caricetosum hostianae* Koch 1926 – Switzerland (Koch 1926);
- 22: *Molinietum caeruleae caricetosum paniceae* Koch 1926 – Switzerland (Koch 1926);
- 23: *Molinietum caeruleae caricetosum tomentosae* Koch 1926 – Switzerland (Koch 1926);
- 24: *Carici davallianae-Molinietum caeruleae* Španíková 1978 – Slovenia (Zelnik & Čarni 2008).

Appendix 2

Dates and localities of the relevés in Table 1

- Rel. 1, 5, 6, 9, 13: Campo di Rovere, Rocca di Mezzo (L'Aquila), 03 July 2023
 Rel. 2–4, 14–15, 17: Campo di Rovere, Rocca di Mezzo (L'Aquila), 04 July 2023
 Rel. 7–8: Campo di Rovere, Rocca di Mezzo (L'Aquila), 30 June 2023
 Rel. 10–13: Le Conche, Ovindoli (L'Aquila), 02 July 2024

Appendix 3

Sporadic species.

- Rel. 1: *Scorzoneroideis cichoriaceae* (Ten.) Greuter +. Rel. 5: *Trifolium hybridum* L. subsp. *hybridum* +. Rel. 6: *Gentiana verna* L. subsp. *verna* +.2; *Parnassia palustris* L. subsp. *palustris* +. Rel. 7: *Alchemilla* sp. 1.2; *Hypericum perforatum* L. subsp. *perforatum* +. Rel. 8: *Campanula glomerata* L. 1.2; *Stellaria graminea* L. +.2; *Campanula rapunculus* L. +; *Lophiolepis tenoreana* (Petr.) Del Guacchio, Bureš, Iamónico & P.Caputo +; *Cruciata laevipes* Opiz +; *Saxifraga bulbifera* L. +; *Tragopogon porrifolius* L. +. Rel. 9: *Carex liparocarpos* Gaudin subsp. *liparocarpos* +. Rel. 10: *Danthonia decumbens* (L.) DC. subsp. *decumbens* 1.2; *Agrimonia eupatoria* L. subsp. *eupatoria* +. Rel. 12: *Valeriana stolonifera* Czern. subsp. *angustifolia* Soó +. Rel. 13: *Anthyllis vulneraria* L. subsp. *weldeniana* (Rchb.) Cullen 1.1; *Phragmites australis* (Cav.) Trin. ex Steud. +. Rel. 16: *Mentha aquatica* L. subsp. *aquatica* +.

Table 1 (Tabela 1): *Centaureo jaceae-Molinietum caeruleae* Ciaschetti, Sburlino et Venanzoni *ass. nova*.

Relevé No.	1*	2	3	4	5	6	7	8	9	10	11	12	13	14*	15	16	17	
Coverage (%)	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	
Area (s. m.)	16	16	16	16	16	20	30	30	16	25	20	16	16	25	20	20	25	
<i>Centaureo jaceae-Molinietum caeruleae ass. nova</i>																		
<i>Molinia caerulea</i> (L.) Moench	4.4	4.4	4.4	4.4	4.4	4.4	2.2	3.4	4.4	5.5	5.5	5.5	4.4	4.4	4.4	4.4	4.4	
<i>Centaurea jacea</i> L. subsp. <i>jacea</i>	1.1	1.1	1.1	2.2	1.1	+	2.2	2.3	2.2	1.1	2.2	+	2.2	2.2	2.2	1.1	2.2	
<i>Rhinanthus minor</i> L.	1.1	+		1.1	1.1	1.1	2.2	1.1	1.1	1.1	2.2	1.1	1.1	+	2.2	1.1	1.1	
<i>Sesleria uliginosa</i> Opiz	3.3	3.3	2.2	2.2	1.1		4.4	4.4	3.3				2.2	2.2	1.1	1.1		
<i>Ranunculus polyanthemoides</i> Boreau	+	+	1.1	+	1.1	1.1			+					2.2				
typicum																		
<i>Genista tinctoria</i> L.	1.2	1.1	1.1	1.1	1.1	1.1	+	2.2	1.2								+	
<i>Filipendula vulgaris</i> Moench	+	+	+	1.1	+	+	3.3	2.2	2.2									
<i>Trifolium montanum</i> L. subsp. <i>rupestris</i> (Ten.) Nyman	1.1	1.1	2.2	2.2	+	+			1.1								+	
<i>Lathyrus pannonicus</i> (Jacq.) Garcke subsp. <i>asphodeloides</i> (Gouan) Bässler	1.1	+		+				1.1	1.1	1.1							1.1	
<i>Plantago media</i> L. subsp. <i>media</i>	+					+	2.2	1.1	1.1									
<i>Poa sylvicola</i> Guss.	1.1			+	+			1.1										
<i>Klasea lycopifolia</i> variant																		
<i>Klasea lycopifolia</i> (Vill.) A. Löve & D.Löve							1.2	3.4	1.2									
<i>Thesium humifusum</i> DC.						1.1	2.2	1.2	1.2									
<i>Bromopsis erecta</i> (Huds.) Fourr.							1.2	1.2	+									
<i>Knaulia arvensis</i> (L.) Coult.							1.1	+	+									
<i>Brachypodium genuense</i> (DC.) Roem. & Schult.								2.2	2.2									
<i>Euphorbia gasparinii</i> Boiss. subsp. <i>sarmitica</i> (Fiori) Pignatti							1.2	+										
<i>Trifolium ochroleucon</i> Huds.							1.1	1.2										
<i>Avenula pubescens</i> (Huds.) Dumort.							+	2										
<i>carictosum hostianae</i> subass. nova																		
<i>Carex hostiana</i> DC.											1.2	+		1.2	1.2	+	1.1	
<i>Juncus inflexus</i> L. subsp. <i>inflexus</i>										+	+	+		+	+2	+	+	
<i>Dactylorhiza incarnata</i> (L.) Soó subsp. <i>incarnata</i>													+	1.1	+	+	+	
<i>Juncus articulatus</i> L. subsp. <i>articulatus</i>										+				+2	+	+	+	
<i>Carex davalliana</i> Sm.					1.2									2.2			1.2	
<i>Carex nigra</i> (L.) Reichard subsp. <i>nigra</i>												1.2						

Relevé No.	1*	2	3	4	5	6	7	8	9	10	11	12	13	14*	15	16	17
Molinio/Molinietalia																	
<i>Succisa pratensis</i> Moench	2.2	2.2	1.2	+	2.2	2.2	3.3	2.2	1.1	2.2	3.3	2.2	3.3	3.3	2.2	2.2	2.2
<i>Sanguisorba officinalis</i> L.	+	2.2	2.2	+	1.1	+2	1.1		1.1	2.2	+	+	1.1	1.1	1.1	1.1	1.1
<i>Serratula tinctoria</i> L. subsp. <i>tinctoria</i>	1.1	2.2	3.3	3.3	3.3	2.2	+	2.2	2.2	1.1	2.2	1.1	1.1	2.2	1.2	+	2.2
<i>Gentiana pneumonanthe</i> L. subsp. <i>pneumonanthe</i>	1.1	1.1	+	1.1	+	1.1	1.1	1.1	+	1.1	2.2	1.1	1.1	2.2		+	2.2
<i>Linum catharticum</i> L. s.l.										1.1	1.1	1.1					
<i>Deschampsia cespitosa</i> (L.) P. Beauv. subsp. <i>cespitosa</i>	+2	+2	+		1.2					+	+	+	1.1	1.1	+		+
<i>Carex panicea</i> L.										1.1	1.1	1.1		+			
<i>Bistorta officinalis</i> Delarbre								1.2	+								+
<i>Carex tomentosa</i> L.																	
<i>Equisetum palustre</i> L.						+2											
<i>Gymnadenia conopsea</i> (L.) R. Br.							+										
<i>Valeriana officinalis</i> L. subsp. <i>officinalis</i>								+									
Molinio-Arrhenatheretea																	
<i>Vicia cracca</i> L.	2.2	+	1.1	+	+	1.1	1.1	1.1	2.2	2.2	+	1.1	1.1	1.1	1.1	+	+
<i>Festuca rubra</i> L. subsp. <i>commutata</i> (Gaudin) Markgr.-Dann.	2.2	2.2	2.2	2.2	2.2	2.2	1.2	1.2	1.1	1.1	1.1	1.1	2.2	1.2	2.2	2.2	1.1
<i>Carex flacca</i> Schreb. subsp. <i>flacca</i>	1.1	2.2	2.2	2.2	2.2	1.1	1.1	1.1	2.2	+2	1.1		+	2.2	3.3	+	1.1
<i>Potentilla erecta</i> (L.) Raeusch.	3.3	3.3	2.2	2.2	1.1	2.2	+	2.2	2.2		1.1	1.1	+	1.1	2.3	2.2	2.2
<i>Trifolium pratense</i> L. subsp. <i>pratense</i>	+	1.1	1.1		1.1	+	2.2	1.2	+		1.1	1.1	+	2.3	2.2	1.1	2.2
<i>Lotus corniculatus</i> L. subsp. <i>corniculatus</i>	1.1	1.1	1.1				2.2	2.2	2.2	+	1.1	+	+	2.2	1.1	1.1	1.1
<i>Ranunculus acris</i> L. s.l.					+2		1.1			1.1	1.1	+	+	1.1	1.1	+	1.1
<i>Trifolium repens</i> L.	+		+		+		2.2	2.2						+2	1.2	1.1	+
<i>Cerastium holosteoides</i> Fr.				+	+		1.1	1.1		1.2	+	+	+				1.1
<i>Holcus lanatus</i> L. subsp. <i>lanatus</i>				+	+		1.2			2.2	1.1	1.1	1.1	1.2			
<i>Lolium interruptum</i> (Desf.) Banf, Galasso, Foggi, Kopecký & Ardenghi	+			+2			+			+	+						1.1
<i>Cynosurus cristatus</i> L.					2.2		1.1	1.1					1.1	1.1	1.1	+	
<i>Anthoxanthum odoratum</i> L.					2.2		2.2			1.1	+		+				1.1
<i>Lathyrus pratensis</i> L. subsp. <i>pratensis</i>							+			1.1	1.1	+	+				+
<i>Carex hirta</i> L.							1.1	1.1		1.1	1.1	+	+		1.1		1.1
<i>Poa trivialis</i> L.								2.3		+		+					
<i>Leontodon hispidus</i> L. subsp. <i>hispidus</i>	+			1.2									1.1				
<i>Colchicum lusitanum</i> Brot.																	
<i>Phleum nodosum</i> L.					1.1												1.1
<i>Dactylis glomerata</i> L. subsp. <i>glomerata</i>							+			+			2.2	+			
<i>Rumex acetosa</i> L. subsp. <i>acetosa</i>							+			1.1		+					

Relevé No.	1*	2	3	4	5	6	7	8	9	10	11	12	13	14*	15	16	17
<i>Potentilla reptans</i> L.										+		+	+				
<i>Scorzoneroides autumnalis</i> (L.) Moench				+									1.1				
<i>Hypericum tetrapterum</i> Fr.							+										
Other species																	
<i>Briza media</i> L.	1.1	1.1	+2	1.1	1.1	1.1	2.3	2.2	1.1	2.2	1.1	2.2	1.1	1.1	2.2	1.1	1.1
<i>Galium verum</i> L. subsp. <i>verum</i>			2.2	1.1	+		1.2	2.2	1.1	1.1	1.1	+	2.2			+	
<i>Dactylorhiza maculata</i> (L.) Soó subsp. <i>saccifera</i> (Brongn.) Diklic						+	1.1				1.1		1.1	+	1.1	+	+
<i>Euphrasia stricta</i> D. Wolff ex J. F. Lehm.				+	1.1	+				+	1.1	1.1					
<i>Cirsium acaulon</i> (L.) Scop. subsp. <i>acaulon</i>				+2		+	2.2	+	1.1								
<i>Cerastium brachypetalum</i> Desp. ex Pers. subsp. <i>brachypetalum</i>		+				+	1.1										
<i>Plantago maritima</i> L. subsp. <i>serpentina</i> (All.) Arcang.		+	2.2													+	+
<i>Luzula multiflora</i> (Ehrh.) Lej. subsp. <i>multiflora</i>					+					+		+	1.1				+
<i>Anacamptis morio</i> (L.) R. M. Bateman, Pridgeon & M. W. Chase				+				+									+
<i>Trifolium pratense</i> L. subsp. <i>semipurpureum</i> (Strobl) Pignatti										1.1	1.1		+				
<i>Carex caryophyllea</i> Latour.			+	+													
<i>Bromus commutatus</i> Schrad. s.l.							+	+2									
<i>Daucus carota</i> L. subsp. <i>carota</i>							+			+							
<i>Thalictrum simplex</i> L. subsp. <i>simplex</i>							1.2							1.1			
<i>Narcissus poeticus</i> L.								+	+								
Sporadic species	1	0	0	0	1	2	2	7	1	2	0	1	2	0	0	1	0