

Short communication

Waiting through the ice age: a low altitude record of *Daphne blagayana* near its north-eastern distribution range limit (Danube Gorge, Carpathian Mountains)

Adriana Petruș-Vancea¹ , Amalia-Raluca Dumbravă^{2,3} ,
Severus-Daniel Covaciu-Marcov¹  & Sára Ferentî¹ 

Key words: relict, endemic, Quaternary, forest, habitat, protected area

Ključne besede: reliktni, endemiti, kvartar, gozd, habitat, zavarovano območje

Abstract

Daphne blagayana is a relict laurel forest species with a restricted distribution in southeastern Europe. It is also sporadic in Romania, where it was previously reported only in a few locations in the Carpathians. In 2025, we identified the Balkan *Daphne* in a region where it had not previously been reported, in the Danube Gorge (Iron Gates Natural Park), southwestern Romania. It populates steep slopes, covered with rare *Quercus petraea* forests in the upper basin of the Mraconia Valley. The new record is a connection between the populations from the Southern Carpathians and those from the Balkan Peninsula. During the last glacial maximum, the region likely served as a refugium for *D. blagayana*, from where it advanced along the Romanian Carpathians. The altitude of this population (647 m) fits within a broader context, as other mountain species were recorded at low altitudes in this region. Although *D. blagayana* is an endemic and relict species, it is not protected in Romania; however, its presence in a protected area with difficult access will likely ensure its survival.

Izvilleček

Daphne blagayana je reliktna vrsta gozdov z omejeno razširjenostjo v jugovzhodni Evropi. Tudi v Romuniji se pojavlja posamično, saj je bila doslej zabeležena le na nekaj nahajališčih v Karpatih. To vrsto smo odkrili na območju, kjer doslej še ni bila znana – v soteski Donave (Naravni park Železna vrata), jugozahodna Romunija. Raste na strmih pobočjih, poraslih z redkimi gozdovi gradna (*Quercus petraea*), v zgornjem delu porečja doline Mraconia. Nova lokacija predstavlja povezavo med populacijami v Južnih Karpatih in tistimi na Balkanskem polotoku. V času zadnjega ledeniškega viška je to območje verjetno služilo kot refugij za vrsto *D. blagayana*. Od tod se je vrsta verjetno širila vzdolž romunskih Karpatov. Nadmorska višina te populacije (647 m) se umešča v širši kontekst, saj so bile v tej regiji tudi druge gorske vrste zabeležene na nižjih nadmorskih višinah, kot je običajno. Čeprav je *D. blagayana* endemična in reliktna vrsta, v Romuniji ni zakonsko zaščitena. Kljub temu bo njena prisotnost znotraj zavarovanega območja in težka dostopnost rastišča najverjetneje zagotovila njeno dolgoročno ohranitev.

Corresponding author:
Adriana Petruș-Vancea
E-mail: adriana_vancea@yahoo.com

Received: 20. 1. 2026
Accepted: 4. 6. 2026



¹ Department of Biology, Faculty of Informatics and Sciences, University of Oradea, Oradea, Romania.

² Iron Gates Natural Park, Orșova, Mehedinți County, Romania

³ Doctoral School of Biomedical Sciences, University of Oradea, Oradea, Romania

Introduction

Within the Carpathian Mountains, the Danube Gorge and the Mehedinţi Mountains shelter numerous Habitat directive species and rare species (Halada et al., 2024). Generally, the Romanian Carpathians are a hot spot of endemic and relict plant species (e.g., Hurdu et al., 2012, 2016; Kliment et al., 2016; Breman et al., 2020), a fact also true for the fauna (e.g., Pop et al., 2010; Teodor et al., 2018; Novák et al., 2019; Dénes et al., 2025). Additionally, the Carpathians, particularly the Southern Carpathians, have been identified in recent years as a glacial refugium (e.g., Šrámková-Fuxová et al., 2017; Lachowska-Cierlik et al., 2020; Zając et al., 2020; Kirschner et al., 2023; Triest et al., 2024). One of the most interesting relict plant species in the Southern Carpathians is *Daphne blagayana* Freyer (Meusel, 1968). It is considered a relict of European laurel forests (Meusel, 1968, 1969), with a small distribution range, occurring almost exclusively in southeastern Europe (Brus, 2001). Besides a few records in the Romanian Carpathians, it is found in scattered locations across the Balkan Peninsula, reaching the eastern Alps in the west (Meusel, 1968), where it is present in northeastern Italy (Martini & Poldini, 1990). *Daphne blagayana* is distributed mainly in the Balkan Peninsula (Derganc, 1902), as it is considered a Balkan species (Hein & Lippert, 2001), or an Illyrian plant, which at its north-western distribution range limit descends at lower altitudes in mountains than further south, alongside other species considered glacial relicts (Derganc, 1908). In Slovenia and Croatia, this Illyrian plant is considered a tertiary relict that survived the ice age in a large part of Slovenia (Horvat, 1929). Other authors considered it an Illyrian Carpathian species (Badea & Cretzoiu, 1941). Although the earliest records of *D. blagayana* in Romania were attributed to another congener species, its presence has been certified since the late 1800s in some locations in the Southern Carpathians (see Derganc, 1904). In 1941, its presence was known in Romania in one location in the Apuseni Mountains, and a few more in the Southern Carpathians, to which was added a new record, situated close to a previous one, but only a few individuals were observed at the edge of a *Fagus sylvatica* L. - *Abies alba* Mill. forest (see Badea & Cretzoiu, 1941). Afterwards, no information appeared in the country, as the same distribution records were presented in subsequent reviews (e.g., Lupşa, 1975–1976). The situation is also generally consistent with the information available in electronic databases (*Daphne blagayana* Freyer in GBIF Secretariat, 2023). Three closely related haplotypes are present in Romania, more than in other parts of the species distribution range

(Fišer Pečnikar et al., 2017). However, there is little data on the distribution of this species in Romania, and the data are old (e.g., Badea & Cretzoiu, 1941; Meusel, 1968; Lupşa, 1975–1976). Thus, information on the distribution of this relict is of great importance, as even in areas where it is widespread, it has a patchy distribution (Fišer Pečnikar et al., 2017). Even in Slovenia, where it was described and has been protected for more than 100 years, it remains poorly known, even with respect to its distribution (Omejc & Brus, 2005). Thus, we present a new distribution record of *D. blagayana* in the Romanian Carpathians in a region where its presence was not previously known (e.g., Meusel, 1968; Lupşa, 1975–1976; *Daphne blagayana* Freyer in GBIF Secretariat, 2023).

Material and methods

The study area is located in southwestern Romania, within the Danube Gorge, in the Iron Gates Natural Park (IGNP). It is a region with a rich biodiversity and biogeographic peculiarities (e.g., Paşcovschi, 1956; Covaciu-Marcov et al., 2009; Schneider-Binder, 2016; Petruş-Vancea et al., 2024; Teodor et al., 2025). The study region (the Almăj Mountains) is a low mountainous area (Mândruţ, 2006) with a warmer climate than most of Romania (Stoenescu et al., 1966; Mândruţ, 2006). The mountain's maximum altitude is only 1224 m (Mândruţ, 2006), reached at about 20 km north from the study site. The study was made in the Mraconia River basin, in eastern IGNP. Mraconia is a left side tributary of the Danube in the gorge, with a relatively large hydrographic basin (Tetelea, 2014). It has a narrow valley, with a sinuous course and steep slopes covered with beech and oak forests. Some tributaries in the upper region are narrow and shaded (Teodor et al., 2025).

The fieldwork was conducted in February, March, and November 2025. Thus, at the beginning of 2025, we have made more transects in the basin of the Mraconia Valley by walking along different forest roads and other paths in the region. Initially, the species was observed accidentally during a field investigation focused on other species groups (especially insects). *Daphne blagayana* was directly identified on the field, using the literature (e.g., Martini & Poldini, 1990; Brus, 2001; Ciocârlan, 2009), based on its clear features. We didn't collect the plants for herbarium; we only photographed them and their habitats. Additionally, we collected some information on other plants in the area, including the neighboring forests, but because it was the beginning of spring, only a few plants were visible. A second location was observed in November 2025, again by chance, while making transects for other species groups.

Results

Daphne blagayana (Figure 1) was identified on the Teiului Valley ("linden valley"), a tributary of the Mraconia River, on a steep slope in the basin's upper region (Figure 2). The Balkan Daphne was observed in two locations. At the first site, the species is present in two small, interconnected glades and in the neighboring forest, with the entire site situated at the upper part of a steep slope. In February–March 2025, we observed hundreds of flowering shoots, especially in sun-exposed areas. The species was well represented throughout the entire site, on a south / south-western-oriented slope, with a 30°–35° inclination. This site is dominated by *Rubus plicatus* Weihe & Nees, which exhibited only leafless branches because our study was conducted in early spring. In the glades, *D. blagayana* was well exposed to sunlight, similar to the neighboring, rare, young *Quercus petraea* (Matt.) Liebl. forest (Figure 3). This is likely a secondary forest, a natural recovery resulting from the exploitation of an old forest (occurring at least 35–40 years ago). Besides *Quercus petraea*, *Quercus robur* L., *Corylus avellana* L., and *Fraxinus excelsior* L. were present. Among herbaceous plants, *Festuca* sp. L., *Genista* sp. L., *Carex* sp. L., *Potentilla* sp. L., and numerous *Erythronium dens-canis* L. individuals were observed.

The medium density of the Balkan Daphne was 50–55 flowering shoots/m² on this populated slope. The plants were healthy and turgescient, with a height of 10–20 cm above the 10–15 cm thick layer of leaf litter. The population was well-shaped, but apparently isolated. The loca-

tion is difficult to access because the region has only a few abandoned forest roads, most of which are overgrown with vegetation. On one side, the site is surrounded by a young *Q. petraea* forest interspersed with *F. sylvatica* individuals; this relatively compact, shaded forest is not populated by *D. blagayana*, and the boundary between the Balkan Daphne population and the forest is obvious. On the other side, there is a slightly more open *Q. petraea* forest, with more sunlight (Figure 3). On this side, there is a gradual transition in the density of *D. blagayana* flowering shoots: they become rarer as tree density increases. The population likely expanded from the glades into the *Q. petraea* regeneration. On 20th February 2025, the plants were blooming, despite snow patches on the ground. Numerous flowering shoots remained on 9th March, when we revisited this site (Figure 1). At this site, the lowest altitude, where flowering shoots were recorded, was at 647±3 m (44°38'52" N / 22°12'14" E), while the upper limit was at 667±3 m altitude (44°38'52" N / 22°12'17" E). The second site with *D. blagayana* was observed in November 2025. Here, we observed hundreds of flowering shoots (Figure 1C) in late autumn. They were situated on a different slope, at 500–700 m distance (44°38'27" N / 22°12'29" E) from the previous site. The habitat is similar, as the species also occupies open areas surrounded by *Q. petraea* and *F. sylvatica* forests. The slope had a 20°–30° inclination, the altitude ranged from 650 to 700 meters, and the plants were present in areas where the forest was more open. As in the previous habitat, both old and young trees were present.



Figure 1: *Daphne blagayana* in the Teiului Valley, Iron Gates Natural Park, Romania, in the spring (A and B) and autumn (C) of the year 2025.

Slika 1: *Daphne blagayana* v dolini Teiului, Narodni park Železna vrata, Romunija, spomladi (A in B) in jeseni (C) leta 2025.




Figure 2: Distribution of *Daphne blagayana* in Romania with old records (according to Lupşa 1975–1976) and the new records' area (above). Details of the area where the Balkan Daphne was recorded in the Iron Gates Natural Park, Romania (below).

Slika 2: Razširjenost vrste *Daphne blagayana* v Romuniji s starimi (po Lupşa 1975–1976) in novimi podatki (zgoraj). Podrobnejši prikaz območja v Naravnem parku Železna vrata v Romuniji, kjer je bila najden *Daphne blagayana* (spodaj).



Figure 3: The *Q. petraea* forest where *D. blagayana* was recorded in the Teiului Valley, Iron Gates Natural Park, Romania.

Slika 3: Gozd gradna (*Q. petraea*), kjer so zabeležili vrsto *D. blagayana* v dolini Teiului Valley, Narodni park Železna vrsta, Romunija.

Discussion

The record of *D. blagayana* in the Mraconia River basin is not just a first record for the Danube Gorge area, but for the entire Banat Mountains, as this represents the most south-western distribution record in Romania, compared with the literature (e.g., Badea & Cretzoiu, 1941; Meusel, 1968; Lupşa, 1975–1976; *Daphne blagayana* Freyer in GBIF Secretariat, 2023). It also appears to be the lowest location in Romania, as previous records ranged from 900 m to 1.890 m in altitude (Lupşa, 1975–1976). However, online databases record it at lower altitudes inside a large urban area (Ştefăneşti Argeş town, near Piteşti city, at about 310 m altitude) (*Daphne blagayana* Freyer in GBIF Secretariat, 2023), indicating a location where the species may have been introduced to gardens, as in other cases (e.g., Strgar, 1975; Brus, 2001). Although at its northern distribution range limit, *D. blagayana* occurs at lower altitudes (e.g., Martini & Poldini, 1990; Dakskobler et al., 2011), in Romania, it was previously observed only at

higher altitudes (Lupşa, 1975–1976). In the north-western parts of its distribution range, few individuals were present at altitudes of 630–670 m in the undergrowth of different *F. sylvatica* forests (Dakskobler et al., 2011). In Slovenia, it descends to 440–460 m, altitudes where there are both native and introduced populations (Omejc & Brus, 2005). In Italy, *D. blagayana* was recorded in two locations between 580 and 930 m altitude (Martini & Poldini, 1990). At its north-western distribution limit, it occurs at 1000 m on steep peaks with little winter snow, but because the area was glaciated, its survival and relict status are difficult to accept (Hein & Lippert, 2001). In this case, the presence of the Balkan *Daphne* likely results from a northward expansion into areas with suitable conditions in the Alps after the last glacial maximum (Hein & Lippert, 2001). In Bulgaria, it is considered a rare, typically mountain species (Yurukov & Zhelev, 2001), connected with *Pinus nigra* forests, whose assem-

blages have a relict origin (Tzonev et al., 2018). It appears that in Bulgaria, *D. blagayana* attains the highest altitude within its range (Fišer Pečnikar et al., 2017). Thus, the location within IGNP has not only biogeographic but also ecological significance, as it represents the species' lowest-altitude record in Romania. The new distribution record is located at approximately 60–70 km south-west of the closest previously known location (the upper basin of the Motru River) (Meusel, 1968; Lupşa, 1975–1976; *Daphne blagayana* Freyer in GBIF Secretariat, 2023) and at approximately 4 km from the Danube. Thus, the new record represents a partial connection with populations south of the Danube, that belong to haplotypes different from those in Romania (Fišer Pečnikar et al., 2017). Only subsequent genetic studies will be able to establish if this geographically intermediate population is also genetically intermediate.

Daphne blagayana likely survived in the Mraconia basin during the Quaternary glaciations, just like other species in the Southern Carpathians (e.g., Schmitt & Varga, 2012; Šrámková-Fuxová et al., 2017; Lachowska-Cierlik et al., 2020; Zajac et al., 2020; Kirschner et al., 2023; Triest et al., 2024). The Balkan *Daphne* was not discovered for a long period in the region (Badea & Cretzoiu, 1941; Meusel, 1968; Lupşa, 1975–1976; *Daphne blagayana* Freyer in GBIF Secretariat, 2023), probably due to its difficult access. However, previously numerous studies were made upon plants in IGNP (e.g. Csűrös et al., 1968; Boşcaiu & Resmeriţă, 1969; Pop et al., 1969; Coldea et al., 1970; Dihoru et al., 1973; Schneider-Binder, 2014), as the region is considered well studied from this perspective (Drăgulescu, 2014), although most studies focused on the areas close to the Danube. Even if the region with *D. blagayana* in the Mraconia Valley was not targeted by such studies, human access there is evidenced by past deforestation, the consequences of which are still visible in the forests of the upper valley. At the same time, this dwarf plant, with only 30 cm in height (Martini & Poldini, 1990; Brus, 2001), is probably difficult to observe. It waited a long time, but its biogeographic significance is now obvious. Thus, *D. blagayana* probably survived in the Mraconia Valley during the glacial periods and subsequently expanded along the Southern Carpathians, as at its northwestern range limit in the Alps (Martini & Poldini, 1990; Hein & Lippert, 2001). In this case, the haplotype became even more relevant, as it is possible that this population belongs to a different one that survived in isolation during the glaciations. It was recently suggested that a mountain insect species (*Liparus glabrirostris* Küster, 1849) could survive during the glaciations in the Mraconia Valley (Teodor et al., 2025). In IGNP, there are likely additional *D. blagayana* populations hidden

on remote, difficult-to-access slopes, as evidenced by our November record of another population on a different slope. This confirms the importance of new studies in the region; as for example, in IGNP, *Vaccinium myrtillus* L. was recently identified at low elevations (Petruş-Vancea et al., 2024).

The presence of *D. blagayana* at low altitude, although unusually related to its distribution in Romania (e.g., Lupşa, 1975–1976), is integrated in a wider biogeographic context of the Danube Gorge, where some mountain species (plants, insects, and vertebrates) descend at very low altitudes (e.g., Paşcovschi, 1956; Csűrös et al., 1968; Covaciu-Marcov et al., 2009, 2025; Teodor et al., 2019; Petruş-Vancea et al., 2024). In the Danube Gorge, the Mraconia Valley is notable in this respect, as recently identified relict mountain species with biogeographic peculiarities were found at low altitudes (Dumbravă & Cupşa, 2025; Teodor et al., 2025). This fact was also confirmed for plants, as certain species were identified in this valley at their lowest altitudes in the Carpathians (Schneider-Binder, 2016). Recently, in the Mraconia basin, a mountain insect species (*L. glabrirostris*) was first identified in the IGNP at low altitude (Teodor et al., 2025). Afterward, hoverfly species (Diptera Syrphidae) with similar requirements were identified in the hydrographic basin (Dumbravă & Cupşa, 2025). The presence of *D. blagayana* is even more relevant than that of the insect (Teodor et al., 2025), which was previously reported in the Banat Mountains, whereas the Balkan *Daphne* is the first record in the region. However, for the insect as well, it was suggested that Mraconia Valley represents a connection between populations from the Southern Carpathians and those from the Balkan Peninsula (Teodor et al., 2025), a fact that may also apply to *D. blagayana*. Thus, the factors that acted upon the insect also acted upon *D. blagayana*, as past conditions allowed the survival of more relict species in this region, confirming that the same conditions from the past acted in the same direction for both plants and animals (Petruş-Vancea et al., 2024). Thus, even if the Danube Gorge is one of the warmest regions in Romania (e.g. Stoenescu et al., 1966; Mândruţ, 2006), with numerous southern, Mediterranean species, at their northern distribution limit (e.g. Paşcovschi, 1956; Coldea et al., 1970; Fuhn, 1975; Tăuşan & Teodorescu, 2017; Dumbravă & Ruicănescu, 2023), Mraconia Valley confirms that the region shelters mountain species at low altitudes (e.g. Paşcovschi, 1956; Schneider-Binder, 2016; Petruş-Vancea et al., 2024; Dumbravă & Cupşa, 2025; Teodor et al., 2025), as the Southern Carpathians have acted as a glacial refugium (e.g. Schmitt & Varga, 2012; Šrámková-Fuxová et al., 2017; Lachowska-Cierlik et al., 2020; Zajac et al., 2020; Kirschner et al., 2023;

Triest et al., 2024). The Danube Gorge area is considered a possible corridor and a potential cryptic refugium for plant species such as *Primula vulgaris* Huds. (see Triest et al., 2024). Additionally, in other plant species, recent data have demonstrated the existence of glacial refuges in the Romanian Carpathians (e.g., Šrámková-Fuxová et al., 2017; Kirschner et al., 2023). This fact also holds for less mobile animals, such as snails (Zajac et al., 2020; Korábek et al., 2023), as well as for more mobile vertebrates (e.g., Horreo et al., 2018). Additionally, lowland populations of a butterfly species were considered potential survivors of glacial periods in the Carpathian valleys (Sitar et al., 2025).

The habitats in the Mraconia basin align with the species' requirements, as *D. blagayana* has previously been recorded in sunny habitats between forests and open areas (Meusel, 1968). Although *F. sylvatica* is well represented in the Danube Gorge (Rozyłowicz, 2008; Manolache et al., 2022), as it occurs even at low altitudes near the Danube (see Paşcovschi, 1956), in the Mraconia Valley, *D. blagayana* is found in *Q. petraea* forests. However, *F. sylvatica* occurs at lower elevations than the Balkan Daphne, in wet, cold, and shady valleys, whereas on the steeper, drier slopes, it is replaced by *Q. petraea*, despite the higher altitude. *Daphne blagayana* was often mentioned alongside conifers (e.g., *A. alba*, *Pinus nigra* J.F. Arnold) (Badea & Cretzoiu, 1941; Bergmeier, 1988). It is considered linked with *P. nigra* (Fukarek, 1957), although this connection could be only a result of the substrate preference (Fukarek, 1969). Naturally, in IGNP, there are no coniferous trees with the exception of *P. nigra*, which is extremely rare, as in the region, there are only deciduous forests (Rozyłowicz, 2008; Manolache et al., 2022), even if many mountain species descend to lower altitudes here (e.g., Paşcovschi, 1956; Schneider-Binder, 2016; Petruş-Vancea et al., 2024; Dumbravă & Cupşa, 2025; Teodor et al., 2025). In Serbia and Montenegro, it occurs in *F. sylvatica* forests, although it is considered characteristic of oak and pine forests (Novakovic-Vukovic et al., 2022). It is present in *F. sylvatica* forests in Bulgaria (Tzonev et al., 2006). *Daphne blagayana* prefers open, light forests; thus, it is decreasing with forest successions (Omejc & Brus, 2005), whereas in IGNP, the steep, rocky slopes did not allow the establishment of a mature forest with old trees. At the same time, past deforestation may have benefited the species by leaving open areas. Although the flowering period takes place in April-May, both in Romania (Badea & Cretzoiu, 1941) and generally (Brus, 2001), in the IGNP, the Balkan Daphne was blooming in February-March, probably because of the lower altitude compared to other locations in Romania (Lupşa, 1975–1976) and of the warmer climate (Stoenescu et al., 1966; Mândruş,

2006). Because of such conditions, in the Danube Gorge, also other mountain species present at lower altitudes began reproducing earlier (Teodor et al., 2025) or were active in winter (Covaciu-Marcov et al., 2022). In the IGNP, *D. blagayana* bloomed again in November, just like in the Alps (Hein & Lippert, 2001).

Although, in certain locations it has been suggested that the species was introduced by irresponsible plant enthusiasts (Accetto, 2010), this possibility is ruled out in our case, as the population's surface area is extensive and the number of flowering shoots is very high. At the same time, in the Mraconia basin (and in neighboring areas), there are many similar steep slopes that are very difficult to access and could shelter other Balkan Daphne populations. Nevertheless, their limited accessibility keeps them undiscovered, a pattern previously observed for this species across much of its range, including its north-western limit (Dakskobler et al., 2011). The large surface area of IGNP and the difficulty of accessing certain areas were considered responsible for incomplete distribution of data in other groups as well (Covaciu-Marcov et al., 2025; Dumbravă & Cupşa, 2025). Probably, the rarity of the species led to it not being reconfirmed even in some areas of the Balkan Peninsula, south of Romania (Sočo et al., 2002).

Although *D. blagayana* is considered protected in all countries within its distribution range by some authors (Fišer Pečnikar et al., 2017), it is not mentioned in the environmental legislation in Romania (O.U.G. 57, 2007). This is an unexpected fact, considering its small range (Brus, 2001), scattered distribution, and relict status (Meusel, 1968, 1969). However, this situation is recurring in Romania for other rare and endemic species that are not mentioned in environmental legislation, even though they have biogeographical importance (Covaciu-Marcov & Ferenţi, 2019). The absence of *D. blagayana* from the Romanian environmental legislation (O.U.G. 57, 2007) is even more confusing, as more than 100 years have passed since it was proposed for protection as a nature monument, a rare and important species (Borza, 1924). Fortunately, in IGNP, *D. blagayana* occurs in a large, protected area with rich biodiversity (e.g., Rozyłowicz et al., 2019), which could help ensure its survival despite lacking a conservation status, through habitat protection. The habitat's relative isolation can also contribute to this direction. Thus, the primary threat to the species in IGNP are future forest interventions, as the difficult access deters tourists who could collect the flowers. Nevertheless, in another region of Romania (Vâlcea County), *D. blagayana* is collected by tourists along with other plants (Ploaie & Turnock, 2001), a practice also observed in other regions, thus it has become rare and endangered

(Paulin, 1902; Brus, 2001). Unlike in Romania, in other regions the species has been protected for more than 100 years (see Omejc & Brus, 2005). In conservation actions for *D. blagayana*, evolutionarily significant units should be used, as Romania is already rich in haplotypes (Fišer Pečnikar et al., 2017), and the population from IGNP could represent a distinct evolutionarily significant unit. Ultimately, the identification of *D. blagayana* confirms that the apparently low number of protected and rare species in the Romanian Carpathians, compared with the Western Carpathians, is attributable to the difficulty of the studies due to the larger country surface area relative to the countries of the Western Carpathians (Halada et al., 2024).

Acknowledgments

Our study was made with the support of the IGNP Administration, to which we are grateful.

Research data availability

All data supporting this study are included within the article.

References

- Accetto, M. (2010). *Daphne blagayana* Freyer vrsta na prvem nahajališču v lškem vintgarju domnevno sajena. *Hladnikia*, 25, 45–67.
- Badea, M., & Cretzoiu, P. (1941). Stațiune nouă de *Daphne blagayana* Frey. în România. *Revista Pădurilor*, 53(5), 285–286.
- Bergmeier, E. (1988). Floristic notes on the Kato Olimbos area (NE Thessaly, Greece). *Willdenowia*, 17(1/2), 37–58.
- Borza, A. (1924). Protecțiune naturii în România. *Buletinul de Informații al Grădinii Botanice și al Muzeului Botanic de la Universitatea din Cluj*, 4(1), 1–24.
- Boșcaiu, N., & Resmeriță, I. (1969). Vegetația ierboasă xerofilă de aluviuni din sectorul Eșelnița – Valea Mraconiei al Defileului Dunării. *Studii și Cercetări de Biologie, Seria Botanică*, 21(3), 209–216.
- Breman, E., Hurdu, B.-I., Kliment, J., Kobiv, Y., Kučera, J., Mráz, P., Pușcaș, M., Renaud, J., Ronikier, M., Šibík, J., Schmotzer, A., Štubňová, E., Szatmari, P.-M., Tasenkevich, L., Turis, P., & Slovák, M. (2020). Conserving the endemic flora of the Carpathian Region: an international project to increase share knowledge on the distribution, evolution and taxonomy of Carpathian endemics and to conserve endangered species. *Plants Systematics and Evolution*, 306, 59. <https://doi.org/10.1007/s00606-020-01685-5>
- Brus, R. (2001). *Daphne blagayana* Freyer. In R. Schütt (Ed.), *Enzyklopädie der Holzgewächse: Handbuch und Atlas der Dendrologie* (pp.1–6). Edition 26. Ergänzungslieferung 12/1.
- Ciocârlan, V. (2009). *Flora ilustrată a României: Pteridophyta et Spermatophyta*. Editura Ceres, București.
- Coldea, G., Boșcaiu, N., Lupșa, V., Plămadă, E., & Resmeriță, I. (1970). Vegetația făgetelor din sectorul Valea Eșelnița – Valea Mraconia al Defileului Dunării. *Studii și Cercetări de Biologie, Seria Botanică*, 22(6), 467–474.
- Covaciu-Marcov, S.-D., & Ferenți, S. (2019). An endemic species in a protected area: *Euscorpius carpathicus* (L., 1767) in the Cozia National Park, Romania (Scorpiones: Euscorpiidae). *Euscorpius*, 279, 1–6. <https://dx.doi.org/10.18590/euscorpius.2019.vol2019.iss279.1>
- Covaciu-Marcov, S.-D., Cicort-Lucaciu, A.-Ș., Gaceu, O., Sas, I., Ferenți, S., & Bogdan, H.V. (2009). The herpetofauna of the south-western part of Mehedinți County, Romania. *North Western Journal of Zoology*, 5(1), 142–164.
- Covaciu-Marcov, S.-D., Lucaciu, B.-I., Maier, A.-R.-M., Cadar, A.-M., Ile, G.-A., Dumbravă, A.-R., & Ferenți, S. (2022). Beyond the victim number: faunistic and ecological data from a road-mortality study in the Iron Gates Natural Park, Romania. *eco.mont – Journal on Protected Mountain Areas Research and Management*, 14(1), 4–13. <https://doi.org/10.1553/eco.mont-14-1s4>
- Covaciu-Marcov, S.-D., Dumbravă, A.-R., & Ferenți, S. (2025). Searching for the missing news: notes on newt distribution in the Iron Gates Natural Park, Romania. *eco.mont – Journal on Protected Mountain Areas Research and Management*, 17(1), 17–24. <https://doi.org/10.1553/eco.mont-17-1s17>
- Csűrös, Ș., Pop, I., Hodișan, I., & Csűrös-Káptalan, M. (1968). Cercetări floristice și de vegetație între Orșova și Eșelnița. *Contribuții Botanice*, 8, 277–312.
- Dakskobler, I., Vončina, A., & Gantar, T. (2011). Rastišča in združbene razmere vrste *Daphne blagayana* v povodju Idrijce. *Hladnikia*, 28, 3–16.
- Daphne blagayana* Freyer in GBIF Secretariat (2023). *GBIF Backbone Taxonomy*. Checklist dataset <https://doi.org/10.15468/39omei> accessed via GBIF.org on 2025-12-18.
- Dénes, A., Dénes, A.-L., & Keresztes, L. (2025). DNA barcode data for the Carpathian headwaters: species-level identification of stonefly (Insecta, Plecoptera) larvae in a biodiversity hotspot of the Apuseni Mountains. *North-Western Journal of Zoology*, 21(1), 20–30.
- Derganc, L. (1902). Geographische Verbreitung der *Daphne blagayana* Freyer. *Allgemeine botanische Zeitschrift für Systematik, Floristik, Pflanzengeographie*, 8, 195–197.
- Derganc, L. (1904). Nachtrag zum Aufsätze über die geographische Verbreitung der *Daphne Blagayana* Freyer. *Allgemeine botanische Zeitschrift für Systematik, Floristik, Pflanzengeographie*, 10, 44–47.
- Derganc, L. (1908). Zweiter Nachtrag zu meinem Aufsätze über die geographische Verbreitung der *Daphne Blagayana* Freyer. *Allgemeine botanische Zeitschrift für Systematik, Floristik, Pflanzengeographie*, 14, 23–24.
- Dihoru, G., Cristurean, I., & Andrei, M. (1973). Vegetația dintre valea Mraconiei – Depresiunea Dubova din Defileul Dunării. *Acta Botanica Horti Bucurestiensis*, 11, 353–423.
- Drăgulescu, C. (2014). The current state of phyto-coenological research in the "Iron Gates" Danube Gorge (Banat, Romania). *Transylvanian Review of Systematical and Ecological Research 16 – special issue, The "Iron Gates" Natural Park*, 59–64. <https://doi.org/10.1515/trser-2015-0033>
- Dumbravă, A.-R., & Ruicănescu, A. (2023). Large, but hard to see: first record of a large European hoverfly *Milesia crabroniformis*

(Diptera, Syrphidae) in Romania. *South-Western Journal of Horticulture, Biology and Environment*, 14(1), 51–56.

Dumbravă, A.-R., & Cupșa, D. (2025). Hoverflies (Diptera, Syrphidae) in the Danube Gorge, Carpathian Mountains, Romania: zoogeographical, ecological and conservation significance. *Acta Zoologica Bulgarica*, 77(4), 545–562. <https://doi.org/10.71424/azb77.4.002930>

Fišer Pečnikar, Ž., Fujs, N., Brus, R., Ballian, D., & Bužan, E. (2017). Insights into the plastid diversity of *Daphne blagayana* Freyer (Thymelaeaceae). *Journal of Systematics and Evolution*, 55(5), 437–445. <https://doi.org/10.1111/jse.12245>

Fuhn, I.E. (1975). Amphibia și Reptilia. *Grupul de Cercetări Complexe Porțile de Fier, seria monografică, Fauna*, 301–303.

Fukarek, P. (1957). Borika (*Daphne blagayana* Frey.) i njena geografska rasprostranjenost. *Šumarstvo*, 10(11/12), 713–722.

Fukarek, P. (1969). Betrachtungen über einige dem gebiete der Balkanländer und Rumänien gemeinsame baum- und Straucharten. *Revue Roumaine de Biologie, Série de Botanique*, 14(1), 33–46.

Halada, L., Lieskovský, J., & Gajdoš, P. (2024). Hotspot of the Habitat Directive species in the Carpathians. *Biological Conservation*, 293, 110572. <https://doi.org/10.1016/j.biocon.2024.110572>

Hein, H., & Lippert, W. (2001). Ein fund von *Daphne blagayana* Freyer in den Chiemgauer Alpen. *Berichte der Bayerischen Botanischen Gesellschaft zur Erforschung der Flora*, 71, 170–171.

Horreo, J.L., Pelaez, M.L., Suárez, T., Breedveld, M.C., Huelin, B., Surget-Groba, Y., Oksanen, T.A., & Fitze, P.S. (2018). Phylogeography, evolutionary history and effects of glaciations in a species (*Zootoca vivipara*) inhabiting multiple biogeographic regions. *Journal of Biogeography*, 45(7), 1616–1627. <https://doi.org/10.1111/jbi.13349>

Horvat, I. (1929). Rasprostranjenje i prošlost mediteranskih ilirskih i pontskih elementa u flori sjeverne Hrvatske i Slovenije. *Acta Botanica Croatica*, 4(1), 1–34.

Hurdu, B.I., Pușcaș, M., Turtureanu, P.D., Niketić, M., Coldea, G., & Zimmermann, N.E. (2012). Patterns of plant endemism in the Romanian Carpathians (south-eastern Carpathians). *Contribuții Botanice*, 47, 25–38.

Hurdu, B.-I., Escalante, T., Pușcaș, M., Novikoff, A., Barth, L., & Zimmermann, N.E. (2016). Exploring the different facets of plant endemism in the South-Eastern Carpathians: a manifold approach for the determination of biotic elements, centres and areas of endemism. *Biological Journal of the Linnean Society*, 119(3), 649–672. <https://doi.org/10.1111/bij.12902>

Kirschner, P., Závěská, E., Hülber, K., Wessely, J., Willner, W., Schönschetter, P., & Frajman, B. (2023). Evolutionary dynamics *Euphorbia carniolica* suggest a complex Plio-Pleistocene history of understorey species of deciduous forest in southeastern Europe. *Molecular Ecology*, 32, 5350–5368. <https://doi.org/10.1111/mec.17102>

Kliment, J., Turis, P., & Janišová, M. (2016). Taxa of vascular plants endemic to the Carpathian Mts. *Preslia*, 88, 19–76.

Korábek, O., Adamcová, T., Pročková, M., Petrusek, A., Hausdorf, B., & Juříčková, L. (2023). In both directions: expansions of European land snail to the north and south from glacial refugia. *Journal of Biogeography*, 50, 654–668. <https://doi.org/10.1111/jbi.14531>

Lachowska-Cierlik, D., Zając, K., Mazur, M.A., Sikora, A., Kubisz, D., & Kajtoch, Ł. (2020). The origin of isolated populations of the mountain weevil, *Liparus glabrirostris* – the flagship species for riparian habitats. *Journal of Heredity*, 111(4), 357–370. <https://doi.org/10.1093/jhered/esaa018>

Lupșa, V. (1975-1976). Plante ocrotite în Parcul Național Retezat și alte masive din Carpații Românești. *Sargetia. Acta Musei Devensis. Științele Naturii*, 11-12, 95–101.

Manolache, S., Rozyłowicz, L., Pătroescu, M., & Dumbravă, A.R. (2022). Iron Gates Natural Park biodiversity. In L. Rozyłowicz, M. Pătroescu, & M.C. Jiplea (Eds.), *Iron Gates Natural Park Monograph* (pp. 85–125). Editura Culturae Hereditatem. <https://doi.org/10.5281/zenodo.7032466>

Martini, F., & Poldini, L. (1990). *Daphne blagayana* Freyer (Thymelaeaceae), nouva per la flora d'Italia. *Webbia*, 44(2), 295–306. <https://doi.org/10.1080/00837792.1990.10670483>

Mândruț, O. (2006). *Mic Atlas de Geografie a României*. Editura Corint, București.

Meusel, H. (1968). Geobotanische beobachtungen in den Südost-Karpaten. *Archiv für Naturschutz und Landschaftsforschung*, 8, 175–210.

Meusel, H. (1969). Zur Ökogeographischen stellung von *Daphne blagayana*. *Revue Roumaine de Biologie, Série de Botanique*, 14(1), 51–56.

Novaković-Vuković, M., Milosević, R., & Čurović, M. (2022). Diversity of plant species in beech and fir forest on various geological substrates in Serbia and National Park “Biogradska Gora” in Montenegro. *Fresenius Environmental Bulletin*, 31(1), 721–727.

Novák, J., Jablonski, D., & Christophoryová, J. (2019). Contribution to the pseudoscorpion fauna of Transylvania and the Eastern and Southern Carpathians, Romania (Arachnida: Pseudoscorpiones). *North-Western Journal of Zoology*, 15(2), 127–134.

Omejc, J., & Brus, R. (2005). Varstvo blagajevga volčina (*Daphne blagayana* Freyer) v gozdnogospodarskem območju Ljubljana. *Zbornik gozdarstva in lesarstva*, 77, 61–83.

O.U.G. nr. 57/2007 privind regimul ariilor naturale protejate, conservarea habitatelor naturale, a florei și faunei sălbatice [Romanian law].

Paulin, A. (1902). Über die geographische Verbreitung von *Daphne blagayana* Freyer. *Mitteilungen Musealver f Krain*, 15, 95–102.

Pașcovschi, S. (1956). Câteva considerații biogeografice asupra Munților Banatului. *Ocrotirea Naturii*, 2, 111–134.

Petruș-Vancea, A., Cupșa, D., Ferent, S., Dumbravă, A.-R., & Covaciu-Marcov, S.-D. (2024). Low altitude *Vaccinium myrtillus* L. populations in the Eșelnița Valley (Danube Gorge, Carpathian Mountains, Romania). *Journal for Nature Conservation*, 82, 126730. <https://doi.org/10.1016/j.jnc.2024.126730>

Ploaie, G., & Turnock, D. (2001). Public perception of environment in the mountains of Vâlcea County. *Geojournal*, 54, 683–701. <https://doi.org/10.1023/A:1021761516756>

Pop, I., Hodișan, I., & Csűrös, S. (1969). Aspecte de vegetație de pe Valea Eșelnița (Mții Almăjului, Banat). *Contribuții Botanice*, 9, 233–243.

Pop, A.A., Pop, V.V., & Csuzdi, C. (2010). Significance of the Apuseni Mountains (the Carpathians) in the origin and distribution of Central European earthworm fauna. *Zoology in the Middle East*, 51(supplement 2), 89–110. <https://doi.org/10.1080/09397140.2010.10638462>

Rozyłowicz, L. (2008). *Metode de analiză a distribuției areal-geografice a testosei lui Hermann (Testudo hermanni Gmelin, 1789) în România, Studiu de caz: Parcul Natural Porțile de Fier*. Editura Universității din București.

Rozyłowicz, L., Nita, A., Manolache, S., Popescu, V.D., & Hartel, T. (2019). Navigating protected areas networks for improving diffusion of conservation practices. *Journal of Environmental Management*, 230, 413–421. <https://doi.org/10.1016/j.jenvman.2018.09.088>

Schmitt, T., & Varga, Z. (2012). Extra-Mediterranean refugia: the rule and not the exception? *Frontiers in Zoology*, 9, 22. <https://doi.org/10.1186/1742-9994-9-22>

Schneider-Binder, E. (2014). Phytogeographical importance of the mountains along the Danube mountain gap valley and surrounding area. *Transylvanian Review of Systematical and Ecological Research*, 16 – special issue, the “Iron Gates” Natural Park, 11–28. <https://doi.org/10.1515/trser-2015-0030>

Schneider-Binder, E. (2016). Riparian vegetation on the left tributaries of the Danube along the “Clisura” cross valley. *Transylvanian Review of Systematical and Ecological Research*, 18(3), 15–30. <https://doi.org/10.1515/trser-2015-0091>

Sitar, C., Sielezniew, M., Malkiewicz, A., Friks, Z.F., Konvička, M., & Konvickova, H. (2025). Eastern arc of glacial relict species – population genetics of Violet Copper *Lycena helle* butterfly in East-Central Europe. *Insects*, 16, 1202. <https://doi.org/10.3390/insects16121202>

Sočo, I., Nikolić, T., Hršak, V., Jelaska, S.D., & Plazibat, M. (2002). The distribution of the genus *Daphne* L. (Thymelaeaceae) in Medvednica Nature Park, Croatia. *Natura Croatica*, 11(2), 225–236.

Šrámková-Fuxová, G., Záveská, E., Kolár, F., Lučanová, M., Španiel, S., & Marhold, K. (2017). Range-wide genetic structure of *Arabis halleri* (Brassicaceae): a glacial persistence in multiple refugia and origin of the Northern Hemisphere disjunction. *Botanical Journal of the Linnean Society*, 185, 321–342. <https://doi.org/10.1093/botlinnean/box064>

Stoenescu, Ș.M., Șchiop, A., Dica, I., Popescu, E., Patrichi, E., & Țepeș, E. (1966). *Atlasul climatologic al R. S. R.*. F.N. București.

Strgar, V. (1975). O varstvu Blagajevga volčina na Slovenskem. *Varstvo Narave (Nature Conservation)*, 8, 67–70.

Tăușan, I., & Teodorescu, B.E. (2017). Contribution to the knowledge of the ant fauna (Hymenoptera: Formicidae) of the Danube Gorges (Romania). *Romanian Journal of Biology – Zoology*, 62(1-2), 21–32.

Teodor, L.A., Czekes, Z., Podluszányi, A., Milin, V.Ș., & Keresztes, L. (2018). Distribution, morphology and ecology of the Carpathian endemic, relict species *Otiorynchus remotegranulatus* Stierlin, 1891, compared to *O. (Elechrans) chrysosus* Boheman, 1843. *North-Western Journal of Zoology*, 14(1), 117–121.

Teodor, L.A., Ferenc, S., & Covaciu-Marcov, S.D. (2019). Weevils Die in Vain? Understanding Messages from Road-Killed Weevils (Coleoptera: Curculionidae). *Coleopterists Bulletin*, 73(2), 359–368. <https://doi.org/10.1649/0010-065X-73.2.359>

Teodor, L.A., Lazăr, A.-M., Mușet, A.-F., & Dumbravă, A.-R. (2025). *Liparus glabrioris* Küster, 1849 (Coleoptera: Curculionidae) in the Danube Gorge: a low altitude population in south-eastern Europe, zoogeographical and ecological considerations. *South-Western Journal of Horticulture, Biology and Environment*, 16(1), 70–79.

Tetelea, C. (2014). Morphometric analysis to extrapolate geoecological potential of the rivers in the “Iron Gates” Natural Park. *Transylvanian Review of Systematical and Ecological Research*, 16 – special issue, the “Iron Gates” Natural Park, 29–46. <https://doi.org/10.1515/trser-2015-0031>

Triest, L., Van Rossum, F., Sramkó, G., Sierens, T., & Volkova, P. (2024). Over the hills and far away: phylogeography and demographic history of a dispersal restricted primrose (*Primula vulgaris*). *Frontiers in Ecology and Evolution*, 12, 1333726. <https://doi.org/10.3389/fevo.2024.1333726>

Tzonev, R., Dimitrov, M., Chytrý, M., Roussakova, V., Dimova, D., Gussev, C., Pavlov, D., Vulchev, V., Vitkova, A., Gogoushev, G., Nikolov, I., Borisova, D., & Ganeva, A. (2006). Beech forest communities in Bulgaria. *Phytocoenologia*, 36(2), 247–279. <https://doi.org/10.1127/0340-269X/2006/0036-0247>

Tzonev, R., Dimitrov, M., Gussev, C., Vulchev, V., & Nikolov, I. (2018). Classification of the relict forest communities of Palla's Black Pine (*Pinus nigra* subsp. *pallasiana*) in Bulgaria. *Hacquetia*, 17(2), 125–161. <https://doi.org/10.1515/hacq-2017-0011%20>

Zajac, K.S., Pročków, M., Zajac, K., Stec, D., & Lachowska-Cierlik, D. (2020). Phylogeography and potential glacial refugia of terrestrial gastropod *Faustina faustina* (Rossmässler, 1835) (Gastropoda: Eupulmonata: Helicidae) inferred from molecular data and species distribution models. *Organism Diversity & Evolution*, 20, 747–762. <https://doi.org/10.1007/s13127-020-00464-x>

Yurukov, S., & Zhelev, P. (2001). The woody flora of Bulgaria: a review. *Schweizerische Zeitschrift für Forstwesen*, 152(2), 52–60. <https://doi.org/10.3188/szf.2001.0052>