





# Diversity, distribution, and cultural significance of Coccinellidae (Ladybirds) in the Dehradun region of Uttarakhand, India

Shafkat Jabbar Mir<sup>1</sup>, Musheerul Hassan<sup>2</sup> , Muhammad Shoaib Amjad<sup>3</sup> , Ivana Vitasović-Kosić<sup>4</sup> , Mohd Majid Jamali<sup>1</sup> & Rainer W. Bussmann<sup>2,5</sup> 

**Key words:** Coccinellidae, altitudinal distribution, habitat variation, zotherapy

**Ključne besede:** Coccinellidae, višinska razširjenost, variabilnost habitata, zooterapija

## Abstract

This study assesses the diversity, distribution, and cultural significance of Coccinellidae (ladybirds) in Dehradun (India), documenting 32 species from 19 genera, with *Coccinella* as the predominant genus, including *C. septempunctata*, *C. transversalis* and *Cheilomenes sexmaculata*. Species richness was greatest at lower altitudes (300 m), with *Coccinella septempunctata* and *Hippodamia variegata* showing a wide altitudinal distribution and a marked decrease with increasing altitude. Agricultural ecosystems harboured the greatest diversity of Coccinellidae, due to the availability of prey and habitat heterogeneity, while flower gardens, mango orchards and forests harboured fewer species. In particular, *Illeis cincta*, *Halyzia sanscrita* and *Harmonia axyridis* were frequently observed in both agricultural and floral habitats, reflecting their ecological adaptability. We studied also ethnozoological practices of Vann Gujjar and Pahari communities, who use species such as *Coccinella hieroglyphica*, *C. leonina* and *Propylea dissecta* to treat ailments such as asthma, cough and skin diseases. A cross-cultural analysis revealed different preferences between the species, with *Coccinella septempunctata* having the highest cultural value due to its symbolic and medicinal significance.

## Izvilleček

Raziskava ocenjuje raznolikost, razširjenost in kulturni pomen družine Coccinellidae (pikapolonice) v Dehradunu (Indija). Dokumentiranih je bilo 32 vrst iz 19 rodov, pri čemer je rod *Coccinella* najpogostejši z vrstami *C. septempunctata*, *C. transversalis* in *Cheilomenes sexmaculata*. Največja vrstna pestrost je bila zabeležena na nižjih nadmorskih višinah (300 m), pri čemer sta vrsti *Coccinella septempunctata* in *Hippodamia variegata* pokazali največji višinski razpon, vendar z izrazitim upadom ob naraščanju nadmorske višine. Največja raznolikost družine Coccinellidae je bila v kmetijskih ekosistemih zaradi razpoložljivosti plena in heterogenosti habitatov, medtem ko je bilo v cvetličnih vrtovih, nasadih manga in gozdovih manj vrst. Vrste *Illeis cincta*, *Halyzia sanscrita* in *Harmonia axyridis* so bile še posebej pogoste tako na kmetijskih kot vrtnih površinah, kar kaže na njihovo ekološko prilagodljivost. Preučili smo tudi etnozoološke prakse skupnosti Vann Gujjar in Pahari, ki uporabljajo vrste, kot so *Coccinella hieroglyphica*, *C. leonina* in *Propylea dissecta* za zdravljenje astme, kašlja in kožnih bolezni. Medkulturna analiza je razkrila preference glede uporabe posameznih vrst, pri čemer ima *Coccinella septempunctata* največjo kulturno vrednost zaradi svojega simbolnega in medicinskega pomena.

Corresponding author:  
Ivana Vitasović-Kosić  
E-mail: ivitasovic@agr.hr

Received: 10. 1. 2025  
Accepted: 2. 4. 2025



<sup>1</sup> School of Agricultural Sciences, Glocal University Saharanpur, U.P, India

<sup>2</sup> Department of Ethnobotany, Institute of Botany, Ilia State University, Tbilisi, Georgia

<sup>3</sup> Department of Botany, Women University of Azad Jammu & Kashmir Bagh, Pakistan

<sup>4</sup> University of Zagreb Faculty of Agriculture, Division of Horticulture and Landscape Architecture, Department of Agricultural Botany, Zagreb, Croatia

<sup>5</sup> Botany Department, State Museum for Natural History, Karlsruhe, Germany

## Introduction

Differences in altitude are an important factor for biodiversity. They create different ecological zones in which unique flora and fauna develop. In the Himalayas, altitude influences temperature, precipitation and habitat structure, leading to significant differences in species composition and diversity (Bhat et al., 2020). For example, lowland zones (300–500 m) are characterised by warmer temperatures and greater habitat heterogeneity, which favours greater species richness, while mid-altitude zones (500–700 m) have transitional ecosystems with moderate diversity. In contrast, higher altitudes are characterised by harsher climatic conditions and less habitat complexity, resulting in lower species richness (Sharma et al., 2023; Suyal et al., 2021).

Habitat differentiation reinforces these patterns. In Dehradun (India), habitats such as forests, agricultural land, mango plantations and market gardens differ significantly in terms of vegetation structure, microclimate and resource availability, which affect the distribution and abundance of insects. For example, agricultural areas and orchards often have higher insect diversity due to abundant prey and favourable microclimates, whereas forests may have lower diversity due to dense tree canopies and limited understory vegetation (Kunwar et al., 2021). However, urbanisation and habitat fragmentation threaten these ecosystems and lead to a decline in biodiversity, especially for specialised insect species (Hassan et al., 2023). Understanding the interplay between altitude, habitat type and biodiversity is crucial for conservation efforts in this ecologically important region.

The Coccinellidae, commonly known as ladybirds, are a diverse and ecologically important insect family. Nearly 6,000 species have been documented worldwide (Kundoo & Khan, 2017), with 510 species in 79 genera reported in India, including 261 predatory species from 57 genera (Pervez, 2011). Despite this diversity, many Indian species of Coccinellidae are still undescribed, probably due to zoogeographic and climatic differences. Coccinellidae are small beetles (0.8–18 mm) with a wide distribution in habitats such as freshwater, coastal areas, foliage, flowers, roots and decaying material (Honek et al., 2017). Their polymorphic nature makes species identification difficult, but their role in biological control programmes is well documented, especially in the control of pests such as aphids, scale insects and mealybugs (Rondoni et al., 2021). Coccinellidae are important natural predators of agricultural pests, contributing to biological control and reducing the need for chemical pesticides (Obrycki & Kring, 1998). They occupy a key position in food webs, serving as both predators and prey, and their presence is

often an indicator of a healthy ecosystem. Their diversity and abundance are influenced by habitat type, altitude and climatic conditions, making them a valuable subject for ecological studies (Hodek & Honek, 1996). However, habitat loss, climate change and pesticide use are increasingly threatening their populations, emphasising the need for research into their distribution and conservation. There is a crucial gap in understanding the diversity of Coccinellidae in the different altitudes and habitats of the Dehradun region.

Dehradun in northern India is globally recognised for its diverse ecosystems, rich biodiversity and unique ethnic communities. The region is home to indigenous groups such as the Vann Gujjars and Pahari, whose traditional ecological knowledge is still under-researched. The Gujjars, a transhumant pastoralist community, migrate seasonally between the Terai Bhabar and the Bugyals in the Himalayas and subsist on buffalo herding (Hassan et al., 2022a). The Pahari, who are mainly engaged in agriculture, have deeply rooted cultural practises and traditional medicinal knowledge (Hassan et al., 2021). Despite their close relationship with the environment, their ethnobiological knowledge, particularly in relation to insect fauna, has not been systematically documented, representing a significant research gap.

To fill these gaps, the present study aims to:


1. to make a comprehensive assessment of the diversity of Coccinellidae at different altitudes and habitats in the Dehradun region,
2. to document and analyse the ethnobiological knowledge and practices of the Vann Gujjar and Pahari communities in relation to insects.

By integrating ecological and ethnobiological approaches, this study seeks to improve our understanding of the diversity of the Coccinellidae and its cultural significance, while contributing to the conservation of biodiversity and traditional knowledge in the region.

## Materials and Methods

### Study area

The present study was conducted in the administrative district of Dehradun in the Indian state of Uttarakhand (Figure 1). Geographically, the region is nestled between the Lesser Himalayas in the north and the Shiwalik Hills in the south, creating a diverse and ecologically rich landscape. Administratively, Dehradun is divided into 8 tehsils (sub-districts) and covers a total area of about 2000 km<sup>2</sup> (Gairola, 2022). According to the Indian Forest Department, the forested area of the district covers



**Figure 1:** Map of the study area and points (red dots) with sampling sites in Dehradun (Uttarakhand, India).

**Slika 1:** Zemljevid raziskovanega območja in točke (rdeče pike) z lokacijami vzorčenja v regiji z Dehradun (Uttarakhand, India).

1608 km<sup>2</sup> and thus makes up a significant part of the region's land area (Gairola, 2022; Khan, 2013). The climate of Dehradun is classified as humid subtropical and is characterised by pronounced seasonal fluctuations. Winters are cool and dry with temperatures ranging from 3 °C to 20 °C, making them a pleasant time of the year. Summers, on the other hand, are hot and intense with temperatures of up to 42 °C. The monsoon season, which lasts from July to September, brings heavy rainfall to the region and transforms the landscape into a lush, green paradise. These seasonal climatic fluctuations favour rich biodiversity and influence the distribution of species in the different habitats (Malik et al., 2021). The unique geographical and climatic conditions of the region make it an ideal location for biodiversity. The interplay between the Lesser Himalayas, the Shiwalik Hills and the Doon Valley creates a mosaic of habitats, including forests, agricultural land, orchards and urban areas, each harbouring distinct ecological community.

## Sampling of coccinellids

The altitudinal and habitat variations of the family Coccinellidae were studied in five altitudinal zones (300 m, 400 m, 500 m, 600 m and 700 m) in the Dehradun district of Uttarakhand. These altitudinal zones were

categorised into lowland (300–500 m) and mid-mountain (500–700 m) zones based on ecological and topographical features. Two sites were selected at each altitude to ensure a comprehensive survey of habitat diversity in the region. The selected sites are shown on the map (Figure 1). The different habitat types, including forests, mango plantations, flower gardens and agricultural areas, were surveyed for the Coccinellidae family at an altitude of 400 m (Figure 2). The collection period was from April to September 2021, using landing nets for vegetation-dwelling species, tapping arches for trees and shrubs and suction cups for small or sensitive species. Larger beetles were collected by hand, while light traps were used to attract nocturnal species. Visual surveys were conducted to ensure comprehensive sampling of all habitats (Achumi et al., 2012). To ensure systematic sampling, an area of 3 ha was randomly selected in each study area, which was divided into 100 squares of 10×10 m<sup>2</sup>. The collected taxa were transferred to glass vials containing ethyl acetate and brought to the laboratory for identification. For preservation, the specimens were dried in an oven at 60 °C for 72 hours. Identification was based on taxonomic keys (Kapur et al., 1956; Kapur, 1963, 1967; Kuznetsov, 1997, Wang et al., 2015) and grey literature. The nomenclature was confirmed using the Global Biodiversity Information Facility (GBIF, 2022).





**Figure 2:** Different selected habitats: (A) Agricultural areas; (B) Flower gardens; (C) Forests; (D) Mango orchards (Photo: Shafkat Jabbar Mir, 2021).  
**Slika 2:** Različni izabrani habitati: (A) kmetijske površine; (B) cvetlični vrtovi; (C) gozdovi; (D) nasadi manga (foto: Shafkat Jabbar Mir, 2021).

## Ethnozoological data collection

From April to September 2022, we interviewed members of the Vann Gujjar and Pahari ethnic groups to participate in the study. Prior to data collection, an exploratory survey was conducted to assess all options and necessary measures. Informants were selected using the snowball method, and only those who showed potential knowledge of beetles were selected for the final interviews. We adhered to the code of ethics of ISE (ISE, 2022), and with prior verbal consent from informants, we collected endemic knowledge through semi-structured interviews followed by group discussions (Hamid et al., 2021; Hassan et al., 2022). A total of 153 informants were selected, including 63 Gujjar and 90 Pahari (Table 1). All these informants belonged to different age groups. Before data collection, the informants were informed about the ob-

jectives of the present study and were able to understand them. Gujjar and Pahari are the languages spoken by Guj-jari and Pahari people, respectively (Ethnologue, 2022).

## Data Analysis

A chord diagram with the circle package was used to visualise the distribution of species across the respective families and their altitudinal differences (Hamid et al., 2021; Hassan et al., 2022). The association analysis between habitat and Coccinellidae fauna was performed using a chord diagram in OriginPro (Haq et al., 2021). A Venn diagram was created using the Bioinformatics and Evolutionary Genomics tool (available at [http://bioinformatics.psb.ugent.be/cgi-bin/liste/Venn/calculate\\_venn.html](http://bioinformatics.psb.ugent.be/cgi-bin/liste/Venn/calculate_venn.html)).

To quantify species diversity, PAST 4.1 was used to calculate the following diversity indices (Kumar et al., 2022)

**Table 1:** Demographic data of respondents from the Vann Gujar and Pahari communities of Dehradun, India.

**Tabela 1:** Demografski podatki intervjuvancev iz skupnosti Vann Gujar in Pahari iz območja Dehradun (India).

Demographic features	Total Percentage	Ethnic Communities	
		Vann Gujar	Pahari
Respondents	153	63 (41.17%)	90 (58.82%)
Male	87 (56.86%)	29	58
Female	66 (43.13%)	34	32
Original Language		Gujjari	Pahari
Socio linguistic characteristics		Monolingual (Elder community) Bilingual (Youngest community)	Monolingual (Elder community) Bilingual (Youngest community)
Age range (27–75)		27–75	27–75
Profession		Sheppard, Housewives	Farmer, Sheppard, Housewives, Craftsmen, Shopkeeper
Livelihood source		Pastoralism	Agriculture

- Shannon-Wiener index ( $H'$ ): Measures species richness and evenness.
- Simpson's Diversity Index (1-D): Assesses dominance and diversity.
- Margalef's Index (D): Estimates species richness.
- Pielou's Evenness index ( $J'$ ): Assesses the evenness of species distribution.
- Fisher's Alpha ( $\alpha$ ): Provides a measure of species richness.

### Use Value (UV)

To evaluate the proportional value of the species, we employed Use value (UV) indices using the formula provided below (Haq et al., 2021).

$$Uv = \sum Ur/N$$

Ur represents the number of use reports for a particular species, and N reveals the total number of respondents.

## Results and Discussions

### Taxonomic Inventory

A total of 32 species of coccinellids were recorded (Table 2; Figure 3). The documented species were categorised into 19 genera. The distribution of species among genera was very uneven, with most genera containing only one species ( $Y = 0.0035x + 1.8246$ ,  $R^2 = 0.0002$ ). The genus *Coccinella* was the most abundant with five species ( $n = 7$ ), followed by *Harmonia* ( $n = 4$ ) and *Miscraspis* ( $n = 3$ ). Each of the remaining genera contained a single species (Figure 3). It is important to note that varieties of species were identified for the genera *Coccinella* and *Cheilomenes*, which are shown in Figure 3.

The highest occurrence of the genus *Coccinella* can be explained by the fact that species of this genus are common in a variety of habitats, e.g. orchards, vegetable gardens and agricultural fields (Obrycki & Kring, 1998). The

**Table 2:** Zootherapy and cultural uses of the collected Coccinellidae fauna across the Vann Gujar and Pahari ethnic groups in Dehradun, Uttarakhand, India. (\*\* Varieties of the species are written in bold)

**Tabela 2:** Zooterapija in kulturna uporaba vzorčene favne družine Coccinellidae v etničnih skupnostih Vann Gujar in Pahari v območju Dehradun, Uttarakhand, Indija. (\*\* varietete vrst so napisane krepko).

Scientific Name	Local name	Abb	Genus	Zootherapy	Cultural Use	Ethnic Groups	Use Value	Use reports
<i>Adalia decempunctata</i> (Linnaeus, 1758)	Gupillu	Ada.dec	<i>Adalia</i>	-----	Be kept in a pot placed in front of patients suffering from an evil spirit disease, and then both are bewitched for some time	Gujjar	0.1	16
<i>Angelis cardoni</i> (Weise, 1990)	Bhundia	Ang.car	<i>Angelis</i>	-----	Black magic	Gujjar	0.1	16
<i>Brumoides suturalis</i> (Fabricius, 1789)	Pillu	Bru.sat	<i>Brumoides</i>	-----	Enchanted and eaten for 40 days to gain magical powers.	Gujjar	0.1	16


Scientific Name	Local name	Abb	Genus	Zootherapy	Cultural Use	Ethnic Groups	Use Value	Use reports
<i>Cheilomenes sexmaculata</i> (Fabricius, 1781)	Gudbinia	Che.sex	<i>Cheilomenes</i>	Sundried, made into powder, added to honey and with walnuts, to treat infertility in both males and females.	People also believe that the frequent occurrence of this species in the home garden is a sign of good luck.	Pahari	0.11	17
<b><i>Cheilomenes sexmaculata</i> var. <i>undulata</i> (Fabricius, 1781)</b> **	Bahv	Che.und	<i>Cheilomenes</i>	Sundried, made into powder, added to honey and with walnuts, to treat infertility in both males and females.	Dried, powdered and applied topically to treat skin diseases, believed to attract the opposite sex.	Gujjar	0.11	17
<i>Chilocorus nigritus</i> (Fabricius, 1798)	Kala Bhundia	Chi.nig	<i>Chilocorus</i>	The whole body is dried and mixed with spider webs to stop bleeding from minor injuries.	This species is made into garlands together with other beetles.	Gujjar	0.11	18
<i>Coccinella hieroglyphica</i> (Linnaeus, 1759)	Kisari	Coc.hie	<i>Coccinella</i>	The whole body is sundried and consumed with honey to treat cough and asthma.	The beetle is dried and burned to produce smoke, which is used against evil spirits.	Gujjar Pahari	0.26	41
<i>Coccinella leonine</i> (Fabricius, 1775)	Sursui	Coc.leo	<i>Coccinella</i>	The whole body is dried in the sun and eaten with honey to treat asthma.		Gujjar	0.22	35
					They are collected and kept overnight in traditional houses to avoid spiders.	Pahari	0.25	39
<i>Coccinella quinquepunctata</i> (Linnaeus, 1758)	Pissu	Coc.qui	<i>Coccinella</i>	The whole body is dried and consumed with lemon and honey to treat infertility.		Gujjar Pahari	0.18	28
<i>Coccinella septempunctata</i> (Linnaeus, 1758)	Quleela	Coc.sep	<i>Coccinella</i>	The whole body is dried in the sun and eaten with honey to treat asthma.	These are collected and kept overnight in traditional houses to avoid spiders. Because of their seven tips, locals believe that they have the potential to cure various problems with evil spirits.	Gujjar Pahari	0.4	62
<b><i>Coccinella septempunctata</i> var. <i>divaricata</i> (Olivier, 1808)</b> **	Cheechad	Coc.div	<i>Coccinella</i>	-----	Placed in a pot in front of patients suffering spiritual disease and then enchanted for ome time	Gujjar Pahari	0.12	19
<i>Coccinella transversalis</i> (Fabricius, 1781)	Mugari	Coc.tra	<i>Coccinella</i>	Species are dried, made into powder, mixed with turmeric, added with fresh running, and applied on the skin to avoid sunburns.	Considered a friend of local farmers because she hunts pests such as aphids.	Pahari	0.12	19
<i>Coccinella undecimpunctata</i> (Linnaeus, 1758)	Bugi	Coc.axy	<i>Coccinella</i>	Species are dried, made into powder, mixed with turmeric, added with fresh running, and applied on the skin to avoid sunburns.	These are placed on the belly of a pregnant woman, as they are said to produce male children.	Gujjar	0.2	32
<i>Henosepilachna vigintioctopunctata</i> (Fabricius, 1775)	Chotha	Epi.vig	<i>Epilachna</i>	-----	Used for the practice of black magic	Gujjar	0.1	16
<i>Halmus chalybeus</i> (Boisduval, 1835)	Saamp ki moasi	Hal.cha	<i>Halmus</i>	-----	Used for the practice of black magic	Gujjar	0.1	16
<i>Halyzia sanscrita</i> (Mulsant, 1846)	Behoor	Hal.san	<i>Halyzia</i>	The whole body is dried and made into a paste by adding egg white and applied topically to treat skin wrinkles.		Gujjar	0.1	16



Scientific Name	Local name	Abb	Genus	Zootherapy	Cultural Use	Ethnic Groups	Use Value	Use reports
<i>Harmonia axyridis</i> (Pallas, 1773)	Teetad	Har.axy	<i>Harmonia</i>	-----	Used for the practice of black magic	Gujjar	0.1	16
<i>Harmonia conformis</i> (Boisduval, 1835)	Patbijena	Har.con	<i>Harmonia</i>	-----	Used for the practice of black magic	Gujjar	0.1	16
<i>Harmonia dimidiata</i> (Fabricius, 1781)	Doh	Har.dim	<i>Harmonia</i>	-----	Used for the practice of black magic	Gujjar	0.1	16
<i>Harmonia octomaculata</i> (Fabricius, 1781)	Mokada	Har.oct	<i>Harmonia</i>	-----	Dead specimens are placed in freshwater early in the morning for some time, then the same water is used for bathing to improve spirituality.	Gujjar	0.1	16
<i>Harmonia dimidiata</i> (Fabricius, 1775)	Trak Makoda	Hen.vig	<i>Harmonia</i>	-----	Sundried specimens are made in powder and applied on the walls of the home to avoid evil-eye.	Pahari	0.1	16
<i>Hippodamia convergens</i> (Guerin-Meneville, 1842)	Drakcheeda	Hip.con	<i>Hippodamia</i>	-----	Used for the practice of black magic.	Pahari	0.1	16
<i>Hippodamia variegata</i> (Goeze, 1777)	Dora	Hip.var	<i>Hippodamia</i>	-----	Used for the practice of black magic.	Pahari	0.1	16
<i>Illeis cincta</i> (Fabricius, 1798)	Ghun	Ill.cin	<i>Illeis</i>	-----	Used for the practice of black magic.	Gujjar	0.1	16
<i>Megalocaria dilatata</i> (Fabricius, 1775)	Bigu	Meg.dil	<i>Megalocaria</i>	-----	Used for the practice of black magic.	Gujjar	0.1	16
<i>Micraspis discolor</i> (Fabricius, 1798)	Sooree	Mic.dis	<i>Micraspis</i>	The whole insect is dried and mixed with turmeric and the resin from <i>Cedrus deodara</i> (Roxb. ex. D.Don) G.Don to treat wounds.	Used for the practice of black magic.	Gujjar	0.15	23
<i>Micraspis vincta</i> (Gorham, 1895)	Teej	Mis.vin	<i>Micraspis</i>	The whole body is dried and mixed with turmeric and the resin from <i>Cedrus deodara</i> (Roxb. ex. D.Don) G.Don to treat wounds.	Used for the practice of black magic.	Pahari	0.1	16
<i>Micraspis allardi</i> (Mulsant, 1866)	Kalikesari	Mic.all	<i>Micraspis</i>	-----	Used for the practice of black magic.	Gujjar	0.1	16
<i>Oenopia sauzeti</i> (Mulsant, 1866)	Mirgachda	Oen.sau	<i>Oenopia</i>	-----	Used for the practice of black magic.	Gujjar	0.1	16
<i>Propylea dissecta</i> (Mulsant, 1846)	Goguu	Pro.dis	<i>Propylaea</i>	The whole in-sects are dried and mixed with turmeric and olive oil to treat sunburns.	Used for the practice of black magic.	Gujjar	0.11	18
<i>Propylea quatuordecimpunctata</i> (Linnaeus, 1758)	Deeraa	Pro.qua	<i>Propylaea</i>	The whole in-sects are dried and mixed with turmeric and olive oil to treat sunburns.	Used for the practice of black magic.	Pahari	0.1	16
<i>Psyllobora bisoetonotata</i> (Mulsant, 1850)	Keedee	Psy.bis	<i>Psyllobora</i>	-----	Used for the practice of black magic.	Gujjar	0.1	16
<i>Rodolia sexnotata</i> (Mulsant, 1850)	Bambeeree	Rod.sex	<i>Rodolia</i>	-----	Used for the practice of black magic.	Gujjar	0.1	16
<i>Rodolia</i> sp. (Mulsant, 1850)	Beendh	Rod.sp	<i>Rodolia</i>	-----	Used for the practice of black magic.	Gujjar	0.1	16







**Figure 4:** Chord diagram showing the identified insect species at different altitudes of Dehradun (Uttarakhand, India). (“N” is number of species). The full names of the species are given in Table 2.

**Slika 4:** Tetivni diagram prikazuje vrste žuželk na različnih nadmorskih višinah v območju Dehradun (Uttarakhand, India). (“N” je število vrst). Celotna imena vrst so prikazana v Tabeli 2.

evations are summarised in Table 3. We observed the lowest number of individuals (291) at an altitude of 600 m, which we attribute to a combination of environmental, ecological and biological factors. Higher altitudes often have harsher conditions, such as lower temperatures and lower oxygen levels, which can limit the survival and reproduction of many species. In addition, the habitat at 600 m altitude may have a lower carrying capacity and harbour fewer individuals. Despite the low number of individuals, the high evenness (0.9245) shows that the few individuals present are evenly distributed among the species. However, the low species richness (as reflected in the Margalef and Fisher\_alpha indices; Table 3) leads to a

lower overall diversity compared to lower elevations. Future studies could investigate the specific factors limiting population size and species richness at this altitude.

This pattern of decreasing diversity with increasing altitude is consistent with previous studies. For example, Arya & Tamta (2016) reported a decline in the diversity of Coccinellidae at higher altitudes in the Kumaon region of Uttarakhand. Similarly, several studies have documented an overall decline in insect diversity with increasing altitude (Joshi & Sharma, 2008; Joshi et al., 2016; Raghavendra et al., 2022; Schowalter, 2022). In particular, species such as *Coccinella transversalis* and *Hippodamia variegata* were found at all altitudes, demonstrating

**Table 3:** Species diversity at various selected altitudes of Dehradun (Uttarakhand, India).

**Tabela 3:** Vrstna pestrost na različnih nadmorskih višinah na območju Dehradun (Uttarakhand, Indija).

Altitude		300 m	400 m	500 m	600 m	700 m
Individuals		2160	1142	990	291	970
Diversity Indices	Dominance (D)	0.04651	0.05743	0.07067	0.0971	0.1280
	Simpson (1-D)	0.9532	0.9421	0.9290	0.9030	0.8720
	Shannon (H)	3.208	2.976	2.731	2.402	2.124
	Evenness (e^H/S)	0.8531	0.8182	0.8543	0.9245	0.9301
	Margalef (D)	3.641	3.21	2.460	1.930	1.160
	Fisher_alpha (α)	4.731	4.270	3.120	2.502	1.369


their adaptability to different environmental conditions. The distribution of *C. transversalis* at different altitudes in Garhwal district of Uttarakhand was also reported by Joshi et al. (2010). Other studies documenting the distribution of insect species at different altitudes are Gurung & Ponnusamy (2019), Saeed et al. (2016), Sharma et al. (2017) and Sajan et al. (2018).

## Habitat selection

The diversity of Coccinellidae was studied in different habitats at an altitude of 400 metres. The greatest diversity was found in agricultural areas (25 species), followed by flower gardens (19 species), mango orchards (16 species) and forests (8 species). The complete list of species and their assignment to habitats can be found in Figure 5. Agricultural areas (Figure 2) have the highest Coccinellidae diversity, which is due to the abundant prey (e.g. aphids, scale insects), the heterogeneity of the habitat due to different crops and marginal vegetation. The open canopy and favourable microclimate favour their dispersal, while human interventions, such as the introduction of beneficial species, encourage their populations. Taken together, these factors make agricultural areas ideal habitats for Coccinellidae compared to flower gardens, mango orchards or forests. The present results are in line with Obrycki & Kring (1998) and Hodek & Honek (1996).


From the Venn diagram (Figure 6a), *Coccinella quinquepunctata*, *Halmus chalybeus*, *Harmonia conformis* and *Propylea dissecta* occurred only in agricultural fields, while *Micraspis vincta* and *Rodolia* sp. occurred only in mango orchards, *Coccinella septempunctata* occurred only in forests, and *Hippodamia convergens* occurred only in flower plantations. In India, Coccinellidae are mainly found in agricultural areas, orchards, forests and urban areas (Obrycki & Kring, 1998).

Species such as *Illeis cincta*, *Halyzia sanscrita*, *Coccinella septempunctata*, *C. leonina*, *Chilocorus nigritus*, *Angelis cardoni*, *Megalocaria dilatata* and *Harmonia axyridis* were found in agricultural areas and flower gardens (Figure 6b). The frequent occurrence of these species in both habitats can be attributed to the vegetation, which provides an ideal environment for Coccinellidae due to the abundance of prey, nectar, pollen, shelter and favourable microclimate. In addition, nursery owners in Dehradun cultivate a variety of plant species (e.g. cucumbers, gourds, aubergines, tomatoes and flowers) for commercial purposes, which are later cultivated by local farmers on agricultural land. The distribution and composition of Coccinellidae species in different habitats is influenced by various environmental factors such as vegetation type, temperature and pH. Taheri & Reyes (2015) reported that vegetation composition plays an important role in determining the distribution of Coccinellidae species.



**Figure 5:** Chord diagram showing the documented Coccinellids fauna in different habitats. The full names of the species are given in Table 2.

**Slika 5:** Tetivni diagram prikazuje vrste družine Coccinellidae v različnih habitatih. Celotna imena vrst so prikazana v Tabeli 2.




**Figure 6:** (a) Venn diagram showing the number of unique and common species in different selected habitats (b) Percentage similarity between different habitats based on the occurrence of identified insect species in Dehradun (Uttarakhand, India).

**Figure 6:** (a) Vennov diagram specifičnih in skupnih vrst v različnih izbranih habitatih (b) odstotek podobnosti med različnimi habitatih na osnovi pojavljanja vrst žuželk v območju Dehradun (Uttarakhand, Indija).

## Zootherapy

Zootherapy, the use of animals for medicinal purposes, has a significant cultural and therapeutic value in traditional practises. In this study, we have documented the ethnozoological use of insect fauna by the Gujar and Pahari communities in Uttarakhand. A total of six diseases (minor injuries, cough, asthma, infertility, skin folds

and sunburn) were treated with ten Coccinellidae species (*Chilocorus nigritus*, *Coccinella hieroglyphica*, *C. leonina*, *C. quinquepunctata*, *C. septempunctata*, *C. undecimpunctata*, *Halyzia sanscrita*, *Micraspis discolour*, *Propylea dissecta* and *Cheilomenes sexmaculata*) in the Gujar community. Among these, asthma was the most common ailment treated by three species (*Coccinella septempunctata*, *C. hieroglyphica* and *C. leonina*) (Figure 7a).



**Figure 7:** (a) Chord diagram showing species used in the Gujar community for a variety of diseases; (b) Chord diagram showing species used in the Pahari community for a variety of diseases. The full names of the species are listed in Table 2.

**Slika 7:** (a) tetivni diagram vrst, ki jih uporabljajo v skupnosti Gujar za različne bolezni; (b) tetivni diagram vrst, ki jih uporabljajo v skupnosti Pahari za različne bolezni. Celotna imena vrst so prikazana v Tabeli 2.

In the Pahari community, five ailments (asthma, cough, infertility, sunburn and minor injuries) were treated by six species (*Coccinella transversalis*, *C. undecimpunctata*, *C. hieroglyphica*, *Micraspis vincta*, *Propylea quatuordecimpunctata* and *Cheilomenes sexmaculata*). Asthma was treated by *Coccinella undecimpunctata* and *C. hieroglyphica* (Figure 7b). These findings are in agreement with Hassan et al. (2023) who reported similar ethnozoological practises among ethnic communities in Jammu and Kashmir, India. A comprehensive overview of the ethnic use of the documented species can be found in Table 2.


## Cross-culture

The chord diagram shows the species used by the Gujar and Pahari communities and illustrates the predominance of the Gujar over the Pahari in the use of Coccinellids species for daily cultural practices (Figure 8a). A total of (N=21), 62% species (i.e., *Chilocorus nigritus*, *Coccinella undecimpunctata*, *Halmus chalybeus*, *Halyzia sanscrita*, *Micraspis discolor*, *Propylea dissecta* and *Adalia decempunctata*) are typical of the Gujar, while only (N=8), 23% species (i.e., *Cheilomenes sexmaculata*, *Coccinella transversalis*, *Hippodamia variegata*, *Micraspis discolor*, *Harmonia dimidiata*, *Propylea quatuordecimpunctata*, *Micraspis vincta*, *Hippodamia convergens*) are typical of the Pahari (Figure 8a). The idiosyncratic species use by the Gujar is due

to non-urbanization, total dependence on nature, especially for food and medicine, and other cultural practices.

A cross-cultural comparison of the documented species showed that there was an overlap between the two selected ethnic communities only in five species (*Coccinella hieroglyphica*, *C. leonine*, *C. septempunctata*, *Coccinella quinquepunctata*, and *Coccinella septempunctata*) which constitute 15% of the documented species (Figure 8b). This can be explained by the fact that these species are commonly used in both ethnic communities due to the prevailing cultural myths and lack of education. *Coccinella septempunctata*, for example, is collected and kept overnight in traditional houses to avoid spiders, and because of its seven spots, locals believe it has the potential to cure various psychological diseases and problems with evil spirits. Similarly, *C. hieroglyphica* is dried and burned to produce smoke that is used against evil spirits. A complete assignment of insect species to ethnic groups can be found in Table 2. Mishra & Tiwari (2012) reported on the traditional use of Coccinellidae by tribal communities in Odisha for medicinal purposes.

The use value (UV) was used to assess the relative importance of the documented species to the local communities, with higher UV indicating greater cultural or practical importance. Seven ethnocultural use categories were identified based on traditional practises: protection from evil spirits, good luck charms, garland making, in-



**Figure 8:** (a) Chord diagram showing the species used by the ethnic communities (Gujar, Pahari); (b) Venn diagram showing the number of unique and common species among Gujar and Pahari in Dehradun (Uttarakhand, India). The complete names of the species are given in Table 2.

**Slika 8:** (a) tetivni diagram vrst, ki jih uporabljajo v etničnih skupnostih (Gujar, Pahari); (b) Vennov diagram števila specifičnih in skupnih vrst v skupnostih Gujar in Pahari v območju Dehradun (Uttarakhand, Indija). Celotna imena vrst so prikazana v Tabeli 2.



secticides, sex determination, black magic and promotion of spirituality. A complete list of UV values for the documented species used by the Gujjar and Pahari communities can be found in Table 2. The highest UV value was found for *Coccinella septempunctata* (0.4), followed by *C. hieroglyphica* (0.26) and *C. leonina* (0.25). These results are in agreement with studies from nearby regions, such as Hassan et al. (2023) and Altaf et al. (2020), which also emphasise the cultural importance of Coccinellidae in traditional practises.

## Conclusions

To summarise, the present study found a significant decline in diversity with increasing elevation, with agricultural land proving to be the predominant habitat for the Coccinellidae family. In particular, agricultural land and flower gardens were found to be common habitats for several species, including *Illeis cincta*, *Halyzia sanscrita* and *Coccinella septempunctata*, with the genus *Coccinella* having the upper hand in the region. Remarkably, many of these species have a traditional medicinal value, a practise known as zootherapy, and a deep cultural significance within the Vann Gujjar and Pahari communities. These tiny creatures not only contribute to the intricate balance of our ecosystem but also play a central role in traditional healthcare and cultural rituals. Furthermore, research into the therapeutic potential of these species could yield new medicinal compounds, offering exciting prospects in the field of biodiversity research and potentially leading to a new economic and cultural recognition of these seemingly inconspicuous organisms. In addition, it is important to conduct further cultural studies to understand the role of coccinellids in the lives of local ethnic communities, which play a central role in maintaining the delicate balance of the local ecosystem.

## Acknowledgements

The authors are grateful to the Gujjar and Pahari communities for disclosing the important ethnozoological knowledge and thank the esteemed reviewers who contributed significantly to improve the quality of the manuscript.

**Funding:** This study is conducted on a purely self-funded basis.




**Author Contributions:** Conceptualization: MH; Methodology: MH, MSA; Data collection: SJM; Data analysis: MH; Supervision: MH, MSA; Initial draft: SJM; Reviewing and revision: MSA, IVK, MMJ, RWB.

**Institutional Review Board Statement:** This research is based on field surveys; no human or human subject's experiments were conducted. The ethical code of the International Society of Ethnobiology was followed (<https://www.ethnobiology.net/what-we-do/core-programs/ise-ethics-program/code-of-ethics/>).

**Informed Consent Statement:** Informed consent was obtained from all informants from the Pahari and Gujjar communities involved in the study.

**Conflicts of Interest:** The authors declare no conflict of interest.

## ORCID iDs

Musheerul Hassan  <https://orcid.org/0000-0001-9884-0945>  
Muhammad Shoaib Amjad  <https://orcid.org/0000-0002-4854-4635>  
Ivana Vitasović-Kosić  <https://orcid.org/0000-0001-9372-5892>  
Rainer W. Bussmann  <https://orcid.org/0000-0002-3524-5273>

## References

- Achumi, B., Hegde, S.N., Lal, P., & Chandra Yeniseti, S. (2013). Altitudinal and seasonal variation in *Drosophila* species on Mount Japfu of Nagaland, a sub-Himalayan hilly state of India. *Journal of Insect Science* 13(1), 117. <https://doi.org/10.1673/031.013.11701>
- Altaf, M., Abbasi, A.M., Umair, M., Amjad, M.S., Irshad, K., & Khan, A.M. (2020). The use of fish and herptiles in traditional folk therapies in three districts of Chenab riverine area in Punjab, Pakistan. *Journal of Ethnobiology and Ethnomedicine* 16(1), 1–21. <https://doi.org/10.1186/s13002-020-00388-y>
- Arya, M.K., & Tamta, P. (2016). Study on distribution and diversity of beetles (Insecta: Coleoptera) in different elevational zones of Binsar Wildlife Sanctuary, Almora, Uttarakhand. *Indian Journal of Entomology and Zoology Studies* 4(4), 311–316.
- Arya, M.K., Tamta, P., & Verma, A. (2018). Systematic survey on alpha diversity of anthophilous insect fauna in Binsar Wildlife Sanctuary, Western Himalaya. *Entomon* 43(2), 99–110.
- Bhat, J.A., Kumar, M., Negi, A.K., Todaria, N.P., Malik, Z.A., Pala, N.A., Kumar, A., & Shukla, G. (2020). Species diversity of woody vegetation along altitudinal gradient of the Western Himalayas. *Global Ecology and Conservation* 24, e01302. <https://doi.org/10.1016/j.gecco.2020.e01302>
- Ethnologue (2022). Languages of the World. Retrieved from <https://www.ethnologue.com/>
- Gairola, S.U. (2022). Traditional medicinal knowledge of plant species found in Maldeota Village, Dehradun, Uttarakhand (India). *Journal of Ethnopharmacology* 285, 114823. <https://doi.org/10.1016/j.jep.2021.114823>

- Goswami, D., Arya, D., & Kaushal, B.R. (2023). Insect diversity in an agroecosystem of Bhabar region of Uttarakhand. *Indian Journal of Entomology* 85(3), 372–380.
- Gurung, B., Ponnusamy, N., & Pal, S. (2019). Species diversity of predaceous coccinellids in different crop ecosystems under the hilly and terai region of West Bengal (India). *Ecology, Environment and Conservation* 25(2), 152–158.
- Hamid, S., Altaf, M., Bussmann, R.W., & Altaf, M. (2021). The ethnic diversities in animal-human interactions in former Jammu and Kashmir State-India. *Ethnobotany Research & Applications* 22(5), 1–15.
- Haq, S.M., Hassan, M., Jan, H.A., Al-Ghamdi, A.A., Ahmad, K., & Abbasi, A.M. (2022). Traditions for future cross-national food security—Food and foraging practices among different native communities in the Western Himalayas. *Biology* 11(3), 455. <https://doi.org/10.3390/biology11030455>
- Haq, S.M., Yaqoob, U., Calixto, E.S., Rahman, I.U., Hashem, A., Abd Allah, E.F., Alakeel, M.A., Alqarawi, A.A., Abdalla, M., Hassan, M., & Bussmann, R.W. (2021). Plant resources utilization among different ethnic groups of Ladakh in Trans-Himalayan Region. *Biology* 10(9), 827. <https://doi.org/10.3390/biology10090827>
- Hassan, M., Haq, S.M., Ahmad, R., Majeed, M., Sahito, H.A., Shirani, M., Mubeen, I., Aziz, M.A., Pieroni, A., Bussmann, R.W., & Yessoufou, K. (2022a). Traditional use of wild and domestic fauna among different ethnic groups in the Western Himalayas – A cross-cultural analysis. *Animals* 12(17), 2276. <https://doi.org/10.3390/ani12172276>
- Hassan, M., Haq, S.M., Amjad, M.S., Ahmad, R., Bussmann, R.W., & Pérez de la Lastra, J.M. (2023). Invertebrates and herptiles for livelihoods – ethnozoological use among different ethnic communities in Jammu and Kashmir (Indian Himalayas). *Frontiers in Pharmacology* 13, 1043155. <https://doi.org/10.3389/fphar.2022.1043155>
- Hassan, M., Yaqoob, U., Haq, M., Lone, F.A., Habib, H., Hamid, S., Jan, H.A., & Bussmann, R.W. (2021). Food and culture: Cultural patterns related to food by indigenous communities in Kashmir—A Western Himalayan region. *Ethnobotany Research and Applications* 22, 1–20.
- Hodek, I., & Honek, A. (1996). *Ecology of Coccinellidae*. Springer.
- Honek, A., Dixon, A.F., Soares, A.O., Skuhrovec, J., & Martinkova, Z. (2017). Spatial and temporal changes in the abundance and composition of ladybird (Coleoptera: Coccinellidae) communities. *Current Opinion in Insect Science* 20, 61–67. <https://doi.org/10.1016/j.cois.2017.04.001>
- International Society of Ethnobiology (ISE) (2022). Code of Ethics. Retrieved from <https://www.ethnobiology.net/what-we-do/core-programmes/ise-ethics-programme/code-of-ethics/>
- Joshi, N.C., Joshi, P.C., Kumar, S., Nath, P., Singh, V.K., & Mansotra, D. (2016). Entomofaunal diversity in fruit orchards along the altitudinal gradients of district Nainital, Uttarakhand (India). *International Journal of Fauna and Biological Studies* 3(5), 113–120.
- Joshi, P.C., & Sharma, P.K. (2008). First records of coccinellid beetles (Coccinellidae) from Haridwar, Uttarakhand, India. *Tropical Natural History* 8(2), 157–167.
- Joshi, P.C., Sharma, P.K., & Khamashon, L. (2010). Species composition and population dynamics of coccinellids from Pauri-Garhwal district of Uttarakhand, India. *Journal of Applied Bioscience* 36(1), 43–47.
- Kapur, A.P. (1956). Systematic and biological notes on the lady-bird beetles predacious on the San Jose Scale in Kashmir with description of a new species (Coleoptera: Coccinellidae). *Records of the Zoological Survey of India* 52(2–4), 257–274.
- Kapur, A.P. (1963). The Coccinellidae of the third Mount Everest expedition, 1924 (Coleoptera). *Records of the Indian Museum* 59(1), 1–12.
- Kapur, A.P. (1967). The Coccinellidae (Coleoptera) of the Andamans. *Proceedings of the National Institute of Sciences of India* 32(3), 148–189.
- Khan, K. (2013). Spatio-temporal analysis of agricultural development: A block-wise study of Dehradun District. *International Journal of Geography and Geology* 2(3), 24–34.
- Kumar, R., Mishra, A., & Singh, S. (2022). Diversity and distribution of Coleoptera in the Western Himalayas: A case study from Uttarakhand, India. *Journal of Insect Conservation* 26(3), 451–463. <https://doi.org/10.1007/s10841-022-00385-4>
- Kundoo, A.A., & Khan, A.A. (2017). Coccinellids as biological control agents of soft-bodied insects: A review. *Journal of Entomology and Zoology Studies* 5(5), 1362–1373.
- Kunwar, R.M., Subedi, B.P., Baral, S.R., Maraseni, T., LeBoa, C., Adhikari, Y.P., & Bussmann, R.W. (2021). Ethnobotany of the Himalayas: Nepal, Bhutanese, and Tibetan Himalayas. In: *Ethnobotany of the Himalayas* (pp. 65–103). Springer. [https://doi.org/10.1007/978-3-030-57408-6\\_5](https://doi.org/10.1007/978-3-030-57408-6_5)
- Kuznetsov, V.N. (1997). *Lady beetles of the Russian Far East*. Memoir No. 1. The Sandhill Crane Press.
- Malik, A., Kumar, A., Kisi, O., Khan, N., Salih, S.Q., & Yaseen, Z.M. (2021). Analysis of dry and wet climate characteristics at Uttarakhand (India) using effective drought index. *Natural Hazards* 105(2), 1643–1662. <https://doi.org/10.1007/s11069-020-04371-4>
- Maqbool, A., Ahmed, I., Kiełtyk, P., & Ceryngier, P. (2018). *Dinocampus coccinellae* (Hymenoptera: Braconidae) utilizes both Coccinellini and Chilocorini (Coleoptera: Coccinellidae: Coccinellinae) as hosts in Kashmir Himalayas. *European Journal of Entomology* 115(1), 1–8.
- Mishra, A.K., & Yousuf, M. (2018). Record of coccinellid beetles (Coleoptera: Coccinellidae) from forest ecosystem of Uttarakhand, India. *Perspectives on Biodiversity of India* 132, 1–10.
- Obyrcki, J.J., & Kring, T.J. (1998). Predaceous Coccinellidae in biological control. *Annual Review of Entomology* 43(1), 295–321. <https://doi.org/10.1146/annurev.ento.43.1.295>
- Pervez, A. (2011). Ecology of aphidophagous ladybird Propylea species: A review. *Journal of Asia-Pacific Entomology* 14(3), 357–365. <https://doi.org/10.1016/j.aspen.2011.01.001>
- Raghavendra, K.V., Bhopaathi, T., Gowthami, R., Keerthi, M.C., Suroshe, S.S., & Ramesh, K.B. (2022). Insects: Biodiversity, threat status, and conservation approaches. *Current Science* 122(12), 1374–1384.
- Rawat, V.K. (2011). *Coccinella septempunctata* (Coccinellidae: Coleoptera) as a predator of cecidozoan, hymenopteran, and dipteran in Garhwal Himalaya, India. *Environment Conservation Journal* 12(3), 59–62.
- Rondoni, G., Borges, I., Collatz, J., Conti, E., Costamagna, A.C., Dumont, F., & Cock, M.J. (2021). Exotic ladybirds for biological control of herbivorous insects—A review. *Entomologia Experimentalis et Applicata* 169(1), 6–27. <https://doi.org/10.1111/eea.12963>

- Saeed, K., Khattak, M.N.K., Naz, F., Akhtar, N., & Khan, M.F. (2016). Genus *Coccinella* (Coccinellidae: Coleoptera) from District Buner, Khyber Pakhtunkhwa, Pakistan. *Entomology, Ornithology & Herpetology* 5(187), 1–5.
- Sajan, K.C., Kapil, K., & Anjali, K. (2018). Records of ladybeetles (Coleoptera: Coccinellidae) from hilly regions of Nepal. *Indian Journal of Entomology* 80(4), 1236–1248.
- Schowalter, T.D. (2022). *Insect ecology: An ecosystem approach*. Academic Press.
- Sharma, P.K., Khanduri, P.C., & Rawat, R.T. (2022). Species richness and abundance of Coccinellids (Coleoptera: Coccinellidae) in agricultural ecosystem of Doiwala region, Dehradun (UK), India. *Environment Conservation Journal* 23(1–2), 395–399.
- Sharma, P.L., Verma, S.C., Chandel, R.S., Chandel, R.P.S., & Thakur, P. (2017). An inventory of the predatory Coccinellidae of Himachal Pradesh, India. *Journal of Entomology and Zoology Studies* 5(6), 2503–2507.
- Sharma, U.C., Datta, M., & Sharma, V. (2023). Climate and biodiversity. In: *Soils in the Hindu Kush Himalayas: Management for agricultural land use* (pp. 27–57). Springer.
- Suyal, D.C., Joshi, D., Kumar, S., Bhatt, P., Narayan, A., Giri, K., Singh, M., Soni, R., Kumar, R., & Yadav, A.N. (2021). Himalayan microbiomes for agro-environmental sustainability: Current perspectives and future challenges. *Microbial Ecology* 82(1), 1–33. <https://doi.org/10.1007/s00248-021-01849-x>
- Taheri, A., & Reyes-López, J.L. (2015). Five new records of ants (Hymenoptera: Formicidae) from Morocco. *Journal of Insect Science* 15(1), 37. <https://doi.org/10.1093/jisesa/iev023>
- Wang, X., Tomaszewska, W., & Ren, S. (2015). A contribution to Asian *Afidentula* Kapur (Coleoptera, Coccinellidae, Epilachnini). *ZooKeys* 516, 35–50. <https://doi.org/10.3897/zookeys.516.9991>