

NOMENCLATURAL NOTES ON SOME ALLIANCES OF THE HALOPHYTIC VEGETATION OF SOUTHERN URAL AND THE CASPIAN LOWLANDS

Tatiana LYSENKO¹ & Ladislav MUCINA^{2, 3}

Abstract

The paper presents validations of some syntaxon names of the saline vegetation of Eastern Europe (southern Russia and Kazakhstan), namely of the *Carici dilutae–Juncion gerardii*, *Anabasio salsae–Artemision pauciflorae*, *Carici dilutae–Juncetum gerardii*. The names of the *Suaedion acuminatae*, *Climacoptero crassae–Suaedion acuminatae* and the *Suaedetum acuminatae* are corrected because of erroneous selection of eponymous taxon names. We also suggest that the ‘*Tamariceto–Salsolion australis*’ *nom. illeg.* should be a synonym of the *Euphorbion sequieranae*.

Keywords: *Artemisietea lerchiana*, *Festuco-Puccinellietea*, *Kalidietea foliati*, *Thero-Salicornietea*.

Izveček

V članku predstavljamo spremembe ali potrditve nekaterih imen sintaksonov slanooljubne vegetacije vzhodne Evrope (južna Rusija in Kazahstan): *Carici dilutae–Juncion gerardii*, *Anabasio salsae–Artemision pauciflorae*, *Carici dilutae–Juncetum gerardii*. Imena *Suaedion acuminatae*, *Climacoptero crassae–Suaedion acuminatae* in *Suaedetum acuminatae* smo popravili zaradi zamenjave imen vrst po katerih so poimenovani sintaksoni. Predlagamo tudi, da se ime ‘*Tamariceto–Salsolion australis*’ *nom. illeg.* uporablja kot sinonim sintaksona *Euphorbion sequieranae*.

Ključne besede: *Artemisietea lerchiana*, *Festuco-Puccinellietea*, *Kalidietea foliati*, *Thero-Salicornietea*.

Abbreviation and nomenclature: ICPN: International Code of Phytosociological Nomenclature, 3rd ed. (Weber et al. 2000); nomenclature of plant species follows EURO+MED PlantBase (2006–).

1. INTRODUCTION

This paper is one of the series of spin-offs of the EuroVegChecklist (Mucina et al., submitted), featuring validation and nomenclature corrections of names of saline and desert vegetation found in southern, continental regions of the Russian Federation and Kazakhstan.

The International Code of Phytosociological Nomenclature (ICPN; Weber et al. 2000) has been applied to perform the validations and corrections. The names of plant taxa follow the Euro+Med PlantBase (www.emplantbase.org).

2. ‘GLAUCO-CARICION DILUTAE’

Golub & Solomakha (1988: 87) have described the *Glauco–Caricion dilutae* to accommodate hygrophilous (sub)saline meadows and pastures steppe and forest-steppe zones of southern Ukraine and Russian Federation. This alliance should be perhaps seen as a Sarmatian biogeographic analogon to the Pannonian *Juncion gerardii* Wendelberger 1943 (see also Vicherek 1973: 80). Golub & Solomakha (1988: 87) have chosen the ‘*Junko–Glaucetum maritimae* Mahn et Schub. 62 em. Klotz et Köck 84’ to serve as the

¹ Department of Phytodiversity Problems, Institute of Ecology of the Volga River Basin RAS, Komzin St 10, Togliatti, 445003, Russian Federation. E-mail: ltm2000@mail.ru

² Iluka Chair in Vegetation Science and Biogeography, School of Plant Biology, The University of Western Australia, 35 Stirling Hwy, Crawley WA 6009, Perth, Australia. E-mail: Laco.Mucina@uwa.edu.au

³ Department of Geography & Environmental Studies, Stellenbosch University, Private Bag X1, Matieland 7602, Stellenbosch, South Africa

holotypus of the *Glauco–Caricion dilutae*. Originally, the ‘*Junco–Glaucetum maritimi* Mahn et Schubert 1962’ was described from Central Germany and Klotz & Köck (1984) misapplied this name for a community from Bashkiria. The *Junco–Glaucetum maritimi* Mahn et Schubert 1962 was considered as synonym of the ‘*Juncetum gerardii* Nordh. 23’ (classified within the *Armerion maritimae* Br.-Bl. et De Leeuw 1926; ‘*Glauco–Puccinellietalia*’, ‘*Asteretea trifolii*’) by Schubert et al. (2001). The Golub & Solomakha’s (1988: 87) typification of the *Glauco–Caricion dilutae* means *de facto* that the latter alliance became a *nomen superfluum* of the *Armerion maritimae*.

The Bashkirian saline vegetation earlier erroneously classified as the *Junco–Glaucetum maritimi* Mahn et Schubert 1962 was later described by Golub et al. (2003: 97) as the ‘*Cariceto secalinae–Scirpetum tabernaemontanii*’ Golub et al. 2003. Besides a need for orthographic correction, this name is invalid due to a technicality – failure to cite the type in *expressis verbis* (ICPN Art. 5, par. 3) as ‘*typus*’ or ‘*holotypus*’. Here we reject the invalid name suggested by Golub et al. (2003: 97) and replace it with valid description of a new association, carrying more appropriate name:

***Carici dilutae–Juncetum gerardii* ass. nov. hoc loco**
Pseudonym (misapplied name): *Junco–Glaucetum maritimi* Mahn et Schubert 1962 *sensu* Klotz & Köck (1984)

Synonym: ‘*Cariceto secalinae–Scirpetum tabernaemontanii*’ Golub et al. 2003 *nom. inval.* (ICPN Art. 5)

Holotypus hoc loco: Klotz & Köck (1984: Table 12, relevé 4)

Diagnostic taxa: *Carex diluta*, *Juncus gerardii*, *Schoenoplectus lacustris* subsp. *glaucus*

Because of the unfortunate typification, the concept of the *Glauco–Caricion dilutae* Golub et Solomakha 1988 became synonymous with the *Armerion maritimae* (see above). We suggest that in Bashkortostan (and possibly in other inland regions of the steppe and forest-steppe zones of southern Russian Federation and Ukraine) the *Juncus gerardii* hygrophilous swards cannot be identified with either the *Armerion maritimae* or the *Juncion gerradii* Wendelberger 1943. We concur with Golub & Solomakha (1988) that this vegetation in Bashkortostan deserves status of an alliance, however, this alliance should receive a new, valid name:

***Carici dilutae–Juncion gerardii* all. nov. hoc loco**
(*Scorzonero–Juncetalia gerardii*, *Festuco–Puccinellietea*)

Pseudonym: *Glauco–Caricion dilutae* Golub et Solomakha 1988

Holotypus hoc loco: *Carici dilutae–Juncetum gerardii* Lysenko et Mucina 2015 (this paper, see above)

Diagnostic taxa: *Agrostis stolonifera*, *Carex diluta*, *Juncus compressus*, *J. gerardii*, *Glaux maritima*

3. ANABASIO SALSAE–ARTEMISION PAUCIFLORAE

Plant communities dominated by *Anabasis salsa* and *Artemisia pauciflora* are supported by solonetz soils of dry and desert steppes and are found also in the northern regions of the desert zone of Russian Federation and Kazakhstan (Karamysheva & Rachkovskaya 1973: 20). The distribution area of this alliance is situated southeast of that of the *Camphorosmo monspeliacae–Artemision pauciflorae* Karpov 2001 *nom. invers. propos.* These communities are classified as a new alliance – the *Anabasio salsae–Artemision pauciflorae*:

***Anabasio salsae–Artemision pauciflorae* Lysenko all. nov. hoc loco**
(*Artemisietalia lerchiana*, *Artemisietea lerchiana*)

Holotypus hoc loco: *Anabasio salsae–Artemisietum pauciflorae* Lysenko 2013 (Lysenko 2013: Tab. 1)

Diagnostic species: *Anabasis aphylla*, *A. salsa*, *Artemisia pauciflora*, *Atriplex cana*, *Leymus ramosus*, *Suaeda physophora*

Photos: Fig. 1B & C

4. ‘TAMARICETO–SALSOLION AUSTRALIS’

The ‘*Tamariceto–Salsolion australis*’ was described by Golub (see Golub & Savchenko 1986; Golub 1994) to accommodate the Caspian desert vegetation on disturbed mobile wind-modelled barchans and raised sand dunes. Habitats supporting the communities of this alliance are found near settlements and farms or on alluvia.

There are problems both with the name as well as the concept of the ‘*Tamariceto–Salsolion australis*’. *Salsola australis* R. Br. was used by Golub (1994) as an eponymous species of the latter




Figure 1: A: A stand of the *Suaedetum acuminatae* on solonchak soils (Russia, Volgograd Region, Kapustin Yar). B: Semi-desert vegetation of the *Anabasio salsae-Artemisietum pauciflorae suaedetosum physophorae* on solonez soils (Russia, Volgograd Region, along eastern and south-eastern lakesides of the Lake Elton). C: The *Anabasio salsae-Artemisietum pauciflorae anabasi-etosum aphyllae* on solonez soils (Russia, Volgograd Region, eastern lakeside of the Lake Elton). Photos: T. Lysenko.

Slika 1: A: Sestoj asociacije *Suaedetum acuminatae* na soločaku (Rusija, regija Volgograd, Kapustin Jar). B: Polpuščavska vegetacija *Anabasio salsae-Artemisietum pauciflorae suaedetosum physophorae* na soloncu (Rusija, regija Volgograd, ob vzhodni in jugovzhodni obali jezera Elton). C: Subasociacija *Anabasio salsae-Artemisietum pauciflorae anabasi-etosum aphyllae* na soloncu (Russia, regija Volgograd, vzhodna obala jezera Elton). Foto: T. Lysenko.

syntaxon. This taxon is considered currently by Uotila (2011) to be synonymous to *Salsola tragus* L. subsp. *tragus* (see also Akhiani et al. 2007). We do not subscribe to this view. *S. australis* R. Br. was described in Australia, and as such it is different from European *S. tragus* subsp. *tragus* (see Borger et al. 2008, Hrusa & Gaskin 2008, Ayers et al. 2009, Chinnock 2010). The taxon reminiscent of *S. australis*, found in the Caspian deserts probably to the *Salsola tragus* s. lat., however, this is pending further studies. In any case, the syntaxon name of the alliance is illegitimate due to taxonomic identification error (ICPN Art. 43). Three associations (*Salsolo australis–Atriplicetum tataricae*, *Salsolo australis–Cannabidentum ruderalis*, *Tamariceto–Salsolietum australis*) classified within the ‘*Tamariceto–Salsolion australis*’ have same illegitimate nomenclatural status (Art. 43).

Due to lingering intricate taxonomy of the *Salsola tragus* complex, we prefer not to correct this name. Secondly, the vegetation described as the ‘*Tamariceto–Salsolion australis*’ can hardly be floristically and ecologically distinguished from the *Euphorbion sequieranae* Golub 1994.

5. ‘*SUAEDION SALSAE*’

In the paper by Freitag et al. (2001: 79) suggested that the authors the name *Suaedion salsae* Golub et Tchorbadze in Golub 1995 (Golub 1995) confused *Suaeda salsa* (L.) Pall. and *Suaeda acuminata* (C. A. Mey.) Moq., and therefore the name has to be corrected:

***Suaedion acuminatae* Golub et Tchorbadze in Golub 1995 corr. Lysenko et Mucina hoc loco**

(*Camphorosmo–Salicornietalia*, *Thero–Salicornietea*)

Corrected name: *Suaedion salsae* Golub et Tchorbadze in Golub 1995 nom. illeg. (ICPN Art. 43)

Synonym: *Suaedion salsae* Golub et Tchorbadze 1988 (ICPN Art. 1)

Holotypus hoc loco: *Suaedetum acuminatae* Golub et Tchorbadze in Golub 1995 corr. Lysenko et Mucina 2015 (this paper)

Diagnostic taxa: *Salicornia perennans*, *Suaeda acuminata*, *S. heterophylla*

The nomenclature type this alliance is the ‘*Suaedetum salsae*’ Golub et Tchorbadze in Golub 1995 that is also illegitimate for the same reason and needs to be corrected to the Article 43 of ICPN:

***Suaedetum acuminatae* Golub et Tchorbadze in Golub 1995 corr. hoc loco**

Corrected name: *Suaedetum salsae* Golub et Tchorbadze in Golub 1995 nom. illeg. (ICPN Art. 43)

Synonym: *Suaedetum salsae* Golub et Tchorbadze 1988 (ICPN Art. 1)

Holotypus hoc loco: Golub & Tchorbadze (1988: Table 11, relevé 6)

Photo: Fig. 1A

6. *CLIMACOPTERO CRASSAE–SUAEDION SALSAE*

The *Climacoptero–Suaedion* was described by Golub & Čorbadze (1989) to accommodate the saline vegetation of Caspian desertic perennial chenopod scrub in wet depressions and alluvia. Golub & Čorbadze (1989) chose the *Kalidietum foliati* to serve as the nomenclature type of the alliance.

One of the eponymous species of the alliance (*Suaeda salsa* (L.) Pall.) was apparently wrongly identified. *Suaeda acuminata* (C. A. Mey.) Moq. Should be seen as the correct identity of this taxon, as also admitted by the senior author of this syntaxonomic concept (see Golub et al. 2001: 92). In the latter publication, a new suballiance concept has been introduced – the *Climacoptero–Suaedion acuminatae* (Golub & Čorbadze 1989; Golub et al. 2001) where *Suaeda acuminata* is rightly used as one of the eponymous species. Yet, this publication fails to introduce the formal correction of the alliance name – an omission that we rectify here:

***Climacoptero crassae–Suaedion acuminatae* Golub et Čorbadze 1989 corr. hoc loco**

(*Kalidietalia caspici*, *Kalidietea foliati*)

Corrected name: *Climacoptero crassae–Suaedion salsae* Golub et Čorbadze 1989 nom. illeg. (ICPN Art. 43)

Phantom: *Climacoptero crassae–Suaedion salsae* Golub et Čorbadze 1988

Holotypus hoc loco: *Kalidietum foliati* Golub et Tchorbadze 1989

7. REFERENCES

- Akhani, H., Edwards, G. & Roalson, E. H. 2007: Diversification of the old world *Salsola* s. l. (Chenopodiaceae): molecular phylogenetic analysis of nuclear and chloroplast data sets and a revised classification. *International Journal of Plant Sciences* 168: 931–956.
- Ayers, D., Ryan, F. J., GrotKopp, E., Bailey, J. & Gaskin, J. 2009: Tumbleweed (*Salsola* section *Kali*) species and speciation in California. *Biological Invasions* 11: 1175–1187. [Published on-line 28 Oct. 2008].
- Borger, C. P. D., Yan, G., Scott, J. K., Walsh, M. J. & Powles, S. B. 2008: *Salsola tragus* or *S. australis* (Chenopodiaceae) in Australia — untangling taxonomic confusion through molecular and cytological analyses. *Australian Journal Botany* 56: 600–608.
- Chinnock, R. J. 2010: Some observations on *Salsola* L. (Chenopodiaceae) in Australia. *Journal of Adelaide Botanical Garden* 24: 75–79.
- EURO+MeD PlantBase 2006–: The information resource for Euro-Mediterranean plant diversity. www.emplabbase.org (last accessed on: 04 November 2014)
- Freitag, H., Golub, V. B. & Yuritsyna, N. A. 2001: Halophytic plant communities in the northern Caspian lowlands: 1, annual halophytic communities. *Phytocoenologia* 31: 63–108.
- Golub, V. B. 1994: The desert vegetation communities of the Lower Volga valley. *Feddes Repertorium* 7–8: 499–515.
- Golub, V. B. 1995: Halophytic, desert and semi-desert plant communities on the territory of the former USSR. Russian Academy of Sciences, Institute of Ecology of the Volga River Basin, Togliatti, 32 pp.
- Golub, V. B. & Tchorbadze, N. B. 1988: K sintaksonomicheskoi kharakteristike rastitel'nykh soobshchestv zapadnykh podstepnykh il'meni del'ty Volgi. (On syntaxonomic characteristic of the plant communities of western sub-steppe Ilmens of the Volga delta). VINITI (Manuscript 6009-B88 08.09.88), Moscow, 57 pp. (in Russian)
- Golub, V. B. & Čorbadze, N. B. 1989: The communities of the order *Halostachyetalia* Ҙopa 1939 in area of western substeppe ilmens of the Volga Delta. *Folia Geobotanica et Phytotaxonomica* 24: 113–130.
- Golub, V. B. & Savchenko, I. V. 1986: Floristicheskaya klassifikatsiya pustynnikh soobshestv del'ty r. Volgi. (Floristic classification of the desert communities the Volga delta river. Syntaxonomy and dynamic of the antropogenic vegetation.) Izd. Bashkirskogo univerziteta, Ufa, pp. 17–26. (in Russian)
- Golub, V. B., Rukhlenko, I. A. & Sokoloff, D. D. 2001: Survey of communities of the class *Salicornietea fruticosae*. *Rastitel'nost' Rossii* 2: 87–98.
- Golub, V. B., Losev, G. A. & Mirkin, B. M. 1991: Aquatic and hygrophytic vegetation of the Lower Volga valley. *Phytocoenologia* 20: 1–63.
- Golub, V. B., Karpov, D. N., Lysenko, T. M. & Bazhanova, N. B. 2003: Conspectus of communities of the class *Scorzonero-juncetea gerardii* Golub et al. 2001 on the territory of the Commonwealth of Independent States and Mongolia. *Byulletin Samarskaya Luka* 13: 88–140.
- Golub, V. B., & Solomakha, V. A. 1988. Vysshie iedinitiy klassifikatsii rastitel'nosti zasolyonykh pochv Evropeiskoi chasti SSSR. (The high-rank units of the classification vegetation on saline soils in the European part of the USSR.) *Byulletin Moskovskogo Obschestva Ispytatelei Prirody, Biologiya* 93: 80–92. (in Russian)
- Hrusa, G. F. & Gaskin, J. F. 2008: The *Salsola tragus* complex in California (Chenopodiaceae): Characterisation and status of *Salsola australis* and the autochthonous allopolyploid *Salsola ryanii* sp. nov. *Madroño* 55: 113–131.
- Karamysheva, Z. V. & Rachkovskaya, E. I. 1973: Botanicheskaya geografiya stepnoi chasti Tsentral'nogo Kazakhstana. (Botanical geography of the steppe region Central Kazakhstan.) Nauka, Leningrad, 279 pp. (in Russian)
- Klotz, S. & Köck, U.-V. 1984: Vergleichende geobotanische Untersuchungen in der Baschkirischen ASSR. 3. Teil: Wasserpflanzen-, Flußufer- und Halophytenvegetation. *Feddes Repertorium* 95: 381–408.
- Lysenko, T. M. 2013: Kharakteristika rastitel'nogo pokrova solontsovykh pochv osobo okhranyemykh prirodnykh territorii – ozer El'ton i Baskunchak. (Characteristic of the vegetation cover on solonetz soils of the protected nature reserves of the Lakes Elton and Baskunchak.) *Vektor nauki TGU* 2(24): 47–53. (in Russian)
- Mucina, L., Bültmann, H., Dierßen, K., Theurillat, J.-P., Raus, T., Čarni, A., Šumberová, K., Willner, W., Dengler, J., Gavilán García, R., Chytrý, M., Hájek, M., Di Pietro, R., Iakush-

- enko D., Pallas, J., Daniēls, F. J. A., Bergmeier, E., Santos Guerra, A., Ermakov, N., Valachovič, M., Schaminée, J. H. J., Lysenko, T., Didukh, Y. P., Pignatti, S., Rodwell, J. S., Capelo, J., Weber, H. E., Solomeshch, A., Dimopoulos, P., Aguiar, C., Hennekens, S. M. & Tichý, L. submitted: Vegetation of Europe: Hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities.
- Schubert, R., Herdam, H., Weinitschke, H. & Frank, J. 2001: Prodrömus der Pflanzengesellschaften Sachsen-Anhalts. Mitteilungen zur Floristischen Kartierung Sachsen-Anhalt, Sonderheft 2: 1–688.
- Vicherek, J. 1973: Die Pflanzengesellschaften der Halophyten- und Subhalophyten vegetation der Tschechoslowakei. Academia, Praha, 200 pp.

Received: 17. 11. 2014

Revision received: 18. 3. 2015

Accepted: 25. 3. 2015